

Fundamentals of Care in Biomechanics— Part 1

Understanding this science is the key to prevention and treatment.

BY MARK MENDESZOON, DPM

Goals and Objectives

After the completion of this CME, the reader will:

- 1) Appreciate the different scientific fields that are applicable to biomechanics
- 2) Recognize the construct of the human musculature and its actions
- 3) Appreciate the three major biomechanical theories applicable to foot and ankle motion
- 4) Understand the phases of the gait cycle
- 5) Differentiate between walking and running biomechanics
- 6) Appreciate anatomy and physiology of the lower extremity
- 7) Recognize and appreciate closed and open chain kinetics
- 8) Understand the role of the mid-tarsal joint and pathomechanics
- 9) Understand the different lower extremity deformities and their impact on overuse injuries
10. Realize the impact of pathomechanics and overuse injuries

Welcome to Podiatry Management's CME Instructional program. Podiatry Management Magazine is approved by the Council on Podiatric Medical Education as a provider of continuing education in podiatric medicine. Podiatry Management Magazine has approved this activity for a maximum of 1.5 continuing education contact hours. This CME activity is free from commercial bias and is under the overall management of Podiatry Management Magazine.

You may enroll: 1) on a per issue basis (at \$27.00 per topic) or 2) per year, for the special rate of \$219 (you save \$51). You may submit the answer sheet, along with the other information requested, via mail, fax, or phone. You can also take this and other exams on the Internet at www.podiatrym.com/cme.

If you correctly answer seventy (70%) of the questions correctly, you will receive a certificate attesting to your earned credits. You will also receive a record of any incorrectly answered questions. If you score less than 70%, you can retake the test at no additional cost. A list of states currently honoring CPME approved credits is listed on pg. 142. Other than those entities currently accepting CPME-approved credit, Podiatry Management cannot guarantee that these CME credits will be acceptable by any state licensing agency, hospital, managed care organization or other entity. PM will, however, use its best efforts to ensure the widest acceptance of this program possible.

This instructional CME program is designed to supplement, NOT replace, existing CME seminars. The goal of this program is to advance the knowledge of practicing podiatrists. We will endeavor to publish high quality manuscripts by noted authors and researchers. If you have any questions or comments about this program, you can write or call us at: **Program Management Services, P.O. Box 490, East Islip, NY 11730, (631) 563-1604 or e-mail us at bblock@podiatrym.com.**

Following this article, an answer sheet and full set of instructions are provided (pg. 142).—**Editor**

Biomechanics is the specialty that applies the laws of mechanics and physics to human performance in order to gain a greater understanding through modeling, simulation, and measurement.¹ Biomechanics requires knowledge in anatomy, kine-

siology, physiology, engineering, and physics, such that the mechanisms of injury are highly understood and thus treatment can be effectively executed.

Anatomical Considerations

Before one can comprehend the magnitude of biomechanics, our

foundation in anatomy and physiology is paramount. The human body has over 600 muscles and over six billion muscle fibers. Of the three types of muscles: cardiac, smooth, and skeletal, our primary focus will be on the skeletal muscle system.

Continued on page 136

Fundamentals (from page 135)

Forty percent of the body's mass is composed of skeletal muscle. Muscle fibers are typically long and cylindrical, measuring 1-40 mm in length and 10-100 microns in diameter.

The functional structure of the human muscle is the sarcomere. It is through the regulation of calcium that muscle contraction is performed (Figure 1). The neuromuscular unit is the area where the nerve cell (motor neuron) meets the muscle fiber (motor end unit). There is only one neuromuscular unit per muscle and it works on an "all or none" principle.²

Muscle macrostructures include muscle, tendons, and ligaments which all have a direct effect on bone, mobility, and function. Muscles have three types of actions (Figure 2). The best-known action is concentric motion in which the fibers shorten and develop significant tension. The best example of this motion is the bicep curl. In exercise, this is known as a positive repetition.

The second action is the eccentric motion in which the muscles elongate and tension is created. The best example of this is running downhill, when the quadriceps must elongate to adjust to picking up too much speed, which could cause injury. This action is known as a braking action or a negative repetition in exercise. This type of muscle activity is the one that causes delayed onset muscle soreness (DOMS). Typically, this occurs 24

hours after exercise.

The last muscle action is the isometric motion, in which the muscle length maintains the same length, and tension develops in time, resisting motion. Holding a dumbbell weight totally straight is a perfect example of this action.³

Tendons

Tendons, also known as sinew, are thick and dense connective tissue that transmits the mechanical force of muscle upon bone. The main cells of the tendons are the fibrocytes. Over 98% percent of tendons are composed of Type 1 collagen. Tendons that change direction in the body typically are protected by a tendon sheath, which has synovial fluid to lubricate the tendon. In the lower extremity, there


Isometric contraction

Muscle contracts but does not shorten




(a)

Concentric contraction



(b)

Eccentric contraction



(c)

Figure 2: Three types of gross muscle actions

The human body has over 600 muscles and over six billion muscle fibers.




Figure 1: Cross section human muscle fiber

are two major tendons that do not have tendon sheaths—the patella and the Achilles tendon. They both have a paratenon to provide some

nourishment and protection to the tendons. Unlike muscles, tendons generally have less than optimal blood supply and thus are at higher risk for injuries and longer recoveries if injured⁴ (Figure 3).

The technical term for tendon pathology is tendinopathy. The term tendonitis is a generalized term used when dealing with in-

juries of the tendons, and this term should be applied to conditions of the tendon only if acute injury or inflammation is involved. Tendinosis is the term utilized for chronic intra-substance disease of the tendon. Tenosynovitis is degeneration of the tendon sheaths. As mentioned before, the patella and Achilles tendon cannot develop tenosynovitis for lack of a tendon sheath.⁵

Ligaments

Ligaments are the dense, fibrous connective tissue that connects bone to bone. Ligaments are comprised of attenuated collagen fibers. These fibers are designed to provide stability to the joints and control forces applied throughout the joints. There are three types of ligaments in the lower extremity: 1) Capsular ligaments are part of the articular capsule; they

Continued on page 137

BIOMECHANICS

Fundamentals (from page 136)

surround synovial joints and provide mechanical reinforcement to the joint. An example is part of the deltoid ligament of the ankle. 2) Extracapsular ligaments, such as the calcaneal fibular ligament, provide joint stability. 3) Intracapsular ligaments allow for more joint motion but generally are weaker ligaments. Examples are the anterior and posterior cruciate ligaments of the knee.⁶

Historical Background

Foot and ankle biomechanics commenced as an interest to physicians in 1935, when Morton discussed anatomic size and hypermobility of the foot and ankle.⁷ In 1942, Manter investigated the subtalar joint and transverse tarsal joint motion.⁸ In 1953, Hicks reviewed normal foot and ankle mechanics, while in 1960, Elfman evaluated the mechanical relationship between the subtalar joint and mid-tarsal joints.⁹

However, in 1954-1966, Dr. Merton Root truly evaluated lower extremity mechanics and defined the normal foot, pathomechanics, and the science behind the fabrication of functional foot orthoses. This was termed foot morphology theory. Root mechanics identifies the subtalar joint position and motion. Ideally, the subtalar joints have three planes of motion, but motion is mainly in the frontal plane. A total of 30 degrees of motion is ideal and thus a 2-1 ratio of inversion to eversion is within normal limits (Figure 4). This has been used as the primary biomechanical foundation in podiatric medical school.¹⁰ Inversion is typically 20 degrees, while eversion is 10 degrees.

In recent decades, two more biomechanics theories of some merit have been explored. Kirby emphasized the tissue stress theory, which relies on the fact

that mechanically-based pathologies of the foot and lower extremity all result from pathological magnitudes of stress acting within the structural components of the foot and lower extremity. Stress is an internal measure of how an object resists a loading force. One measures this by dividing the cross-sectional area of that object by the loading force being applied.¹¹

Classification of Foot Pathology

Classification of foot pathology can be divided into rearfoot varus and valgus as well as forefoot varus and valgus. Rigid or structural rearfoot varus will remain in a fixed position due to alignment of the tibia, calcaneus, or the position of the subtalar joint. A flexible rearfoot varus

The functional structure of the human muscle is the sarcomere.

Dananberg proposed the sagittal plane theory. During walking, the center of body mass must pass from behind the weight-bearing foot to in front of it.¹² As all of these theories have merit, and as biomechanics is not an absolute science, it has been found over the years that utilization of all three theories is in the best interest of patient care.

will be determined by the midtarsal joint. Coleman block testing is very helpful in assessing this type of foot deformity. The direct connection of the rearfoot to the forefoot is via the midtarsal joint. The midtarsal joint consists of the talonavicular joint and the calcanealcuboid joint.

Chopart described this joint in detail, and thus it is known as Chopart's

joint. The midtarsal joint also has triplane motion like the subtalar joint, but it also has its majority of motion in the frontal plane, leading mainly to inversion and eversion of the joint. There are two axes of motion that comprise the midtarsal joint: the longitudinal axis and the oblique axis. The longitudinal axis is parallel with the transverse and sagittal planes. Therefore, motion is only in the frontal plane—inversion and eversion. Due to the unique alignment of the oblique axis, it has minimal motion technically in the sagittal and transverse plane. The average angulation of the midtarsal joint is approximately 55 degrees from the sagittal and transverse planes (Figure 5).

The midtarsal motion is directly related to the subtalar joint's position. As the subtalar joint goes from maximal pronation to supination, the longitudi-




Figure 3: Lower leg anatomy

Continued on page 138

Fundamentals (from page 137)

nal axes of the two midtarsal joints progressively diverge, and thus congruity of the midtarsal joint is lost and motion decreases. This relationship of the subtalar and midtarsal joints provides the ideal standard clinical index position in normal foot mechanics and is known as being ‘locked and loaded’. This position provides the forefoot with all five metatarsal heads being on the same plane and parallel. The calcaneus is perpendicular to the long axis of the tibia and perpendicular to the ground (Figure 6).

Forefoot Varus

Forefoot varus will require the subtalar joint to be in a supinated position. Midtarsal joint motion is diminished, and the forefoot is in a varus position in which the forefoot’s plantar plane is inverted. Typically in this foot type, a significant pressure will be placed underneath the fifth metatarsal.

Forefoot Valgus

Forefoot valgus will require the subtalar joint to be maximally pronated, resulting in an increased range of motion of the midtarsal joint, with the forefoot plane everted. Opposed to a forefoot varus deformity, a valgus deformity will cause significant pressure underneath the first metatarsal.

As motion and forces continue to move forward from the subtalar and midtarsal joints, there are requirements for the first metatarsal phalangeal joint to work effective-

ly. These requirements will necessitate for the subtalar joint to be in a supinated position and for the midtarsal joint to be stable—‘locked and loaded’.¹³ This will allow for the first metatarsal head to plantarflex below the plane of the second metatarsal, therefore engaging the windlass mechanism of the plantar fascia. As the first metatarsal head plantarflexes with forward momentum, the hallux proximal base will glide on the dorsal aspect of the metatarsal head allowing for adequate dorsiflexion. Intrinsic factors such as a normal metatarsal head and center of axis, along with sesamoid and soft tissue pliability, will assist in




Figure 4: Normal subtalar joint motion




Figure 5: Midtarsal joint axes of motion

this mechanism of action. Ideally, approximately eighty plus percent of the body’s weight will move along the first metatarsal phalangeal joint, and the joint should dorsiflex greater than 40 degrees in the ideal foot.

When faulty mechanics ensue via the subtalar joint excessively pronating and the midtarsal joint becoming mobile, the first ray will also have impaired mechanics. Instability of the first ray will be caused by the peroneal longus losing its mechanical advantage and its ability to plantarflex the first ray, causing dorsiflexion of the first ray and affecting normal range of motion of the joint. In turn, the windlass

mechanism weakens or reverses, and this allows for further dorsiflexion of the first ray, which limits motion. The line of progression changes and shifts medially, which causes excessive medially forefoot loading (Figures 7 & 8). Eventually, a hallux limitus or hallux abductovalgus deformity will develop and over time, a cascade of compensatory conditions will ensue, such as lesser metatarsalgia, hammertoes, hyperkeratotic skin conditions, and even arthritis.¹⁴

Pes Plano Valgus

The etiology of pes plano valgus deformity can be derived from genetics, trauma, age, and of course biomechanical factors. Pes plano valgus can be further divided into rigid and flexible classifications. Rigid classifications, which are less prevalent, include congenital vertical talus and tarsal coalitions. Congenital vertical talus generally needs immediate surgical correction in the infant and tarsal

Continued on page 139




Figure 6: Subtalar and midtarsal joint motion relationship

BIOMECHANICS

Fundamentals (from page 138)

coalition typically will be addressed as a teenager due to the secondary ossification centers of the tarsal bones. Regardless, both conditions typically lead towards arthritis with age and thus altered biomechanics.¹⁵

Flexible Flatfoot

Flexible flatfoot deformity is the most common type of foot deformity and is directly related to impaired biomechanics. Understanding planar dominance of deformity is necessary to institute a proper treatment plan for the patient. The three planes of deformity are sagittal plane, frontal plane, and transverse plane. Sagittal plane deformities include posterior leg muscle group equinus, plantarflexion of the talus, medial column breach, and elevated first ray. Frontal plane deformities include calcaneal valgus and everted forefoot. Transverse plane deformity, which is least appreciated, will have an abducted forefoot to rearfoot relationship.

It is imperative to obtain weight-bearing radiographs to determine the planar dominance of deformity. The important anterior to pos-




Figure 7: Normal first ray mechanics



Figure 8: First ray range of motion




Figure 9: Lateral radiograph pes plano valgus

terior radiographic measurements to recognize are that the talo-calcaneal angle will be increased. Talonavicular joint coverage, which typically should be 80-90%, is diminished. A bisection of the talus and first metatarsal should be a straight line that will have the talus bisection line exit medially to the first metatarsal. A resultant angle will form. The calcaneal-cuboid angle typically should be 0 degrees and will have an increased angle. In

The navicular and cuneiform alignment should be parallel, and in a pes planus deformity there will be a 'fault or sag' at the joint level. In some situations, the first ray may be elevated with the metatarsal head dorsiflexed compared to the base of the hallux phalanx base (Figure 9).

A calcaneal axial and sesamoid axial view will evaluate the calcaneal position and plane of the metatarsal heads and determine if they are neutral or in valgus or everted position. As pes plano valgus deformity is progressive, it is necessary to obtain ankle radiographs to evaluate if there is ankle joint deformity. All radiographs should be evaluated for arthritis.¹⁷

Gait Cycle

The human adult walking gait cycle is developed at about the age of 4. The adult cycle requires a single leg stance. The walking gait cycle is comprised of two sections: the swing phase and the contact phase. Although the swing phase has a subtle impact on biomechanics, it is the contact phase which is the leading cause of lower extremity injuries. The contact phase is approximately 62% and the swing phase 38% of

Flexible flatfoot deformity is the most common type of foot deformity and is directly related to impaired biomechanics.

severe cases, a skew foot deformity may present itself and several faults on radiographs will be noted.¹⁶

Sagittal plane radiographs will reveal a low calcaneal inclination angle less than the ideal 20 plus degrees. Plantar flexion of the talus will be noted. Bisection of the talus and first metatarsal should be a straight line and 0 degrees; however, in a flat foot deformity, the talus line will extend below the first metatarsal and create an angle.

the gait cycle. The contact phase has three segments: heel contact, mid-stance, and propulsion. The average time it takes to cycle through the contact phase is about 0.4 seconds. With walking, there is roughly four times the body weight applied with each step and the average person should achieve between 5,000 and 10,000 steps per day with the goal of surpassing the higher number.

Continued on page 140

Fundamentals (from page 139)

Running Gait Cycle

The running gait cycle differs from the walking gait cycle in that there are greater forces applied, a shorter stance phase, longer swing phase, and unlike the walking gait cycle, there is a float phase. This float phase is where the body is actually completely off the ground. Thus with the float phase in running, the body will absorb up to eight times the body weight with each stride. Running one mile could take about an average of 1,500 steps. Regardless of walking or running, it is within the mid-stance phase that the majority of chronic overuse injuries develop (Figure 10).

When evaluating a patient for a biomechanical examination, it is imperative that an open chain or non-weight-bearing and weight-bearing or closed chain examination are performed. Evaluate the patient from head

to toe and investigate for general body relationship and symmetry, muscle flexibility and strength, joint range of motion, leg-length discrepancy. Evaluation of shoes and sock gear is a must. In addition to evaluating the patient lying and sitting down, attempt to eval-

efficiently and effectively. The saying “core to the floor” or “floor to the core” emphasizes the direct relationship of the lower extremity.

Functional screening tests to evaluate core strength include: single leg stands and squats, step down squats,

The majority of overuse conditions develop in the midstance part of the gait cycle.

uate standing and walking as well. A thorough history and physical is paramount in assessing a patient’s condition before actual examination.

Once a general overview of the body is evaluated, a lower extremity focus should start at the core and its relative structures. The core is the link to providing good posture and trunk stability allowing for the back, legs, and feet to function more

star excursion tests (SEBT), swing test, and one in which the patient, sitting crisscross or Indian style, attempts to stand up without leaning towards one side or using hands. **PM**

References

¹ Seigler, Sorin, Foot and ankle joint biomechanics, Sorin Seigler, Journal of Foot and Ankle Biomechanics Vol. 7 s5 2007.

Continued on page 141




Figure 10: Walking & running gait cycle

Fundamentals (from page 140)

² Frontera WR, Ochala J, Skeletal muscle: a brief review of structure and function, *Calcif Tissue Int.*, 2015 Mar;96(3):183-95.

³ Guo, J, et al., Massage Alleviates Delayed Onset Muscle Soreness after Strenuous Exercise: A Systematic Review and Meta-Analysis, *Front Physiol*, 2017 Sep 27;8.

⁴ Binstead, JT, Bhimiji SS, Anatomy, Lower Limb, Calf., StatPearls [Internet]. Treasure Island (FL): 2017.

⁵ McCreesh, Karen, Lewis, Jeremy, Continuum model of tendon pathology—where are we now? *Int J Exp Pathol.* 2013 Aug; 94(4): 242–247.

⁶ Golano Pau, Vega, Jordi et al., Anatomy of the ankle ligaments: a pictorial essay, *Knee Surg Sports Traumatol Arthrosc.* 2010 May; 18(5): 557–569.

⁷ Nester Christopher J, Leardini Alberto, International Foot and Ankle Biomechanics Community (i-FAB): past, present and beyond, *J Foot Ankle Res.* 2009; 2: 19.

⁸ Van Ghelue, Bart, Kirby, Kevin, Research and clinical synergy in foot and lower extremity biomechanics, *Journal Footwear Science*, Vol. 2, 2010.

⁹ Mann, RA, Biomechanics of the Ankle, *Joint Surgery Up to Date* pp 73-81.

¹⁰ Kirby Kevin, Are Root Biomechanics Dying, *Podiatry Today*, vol.22, issue 4, 2009.

¹¹ Kirby, Kevin, Prescribing Orthoses: Has Tissue Stress Theory Supplanted Root Theory?, *Podiatry Today*, vol. 28, issue 4, 2015.

¹² Dananberg Howard, Sagittal Plane Biomechanics, *J Am Podiatr Med Assoc*, 90 (1): 47-50, 2000.

¹³ Blackwood C. Brian, Sangeorzon, The Midtarsal Joint Locking Mechanism, *Foot and Ankle International*, Volume: 26 issue: 12, page(s): 1074-1080, 2005.

¹⁴ Faber Frank, Kleinrensink G.J. et al., Mobility of the First Tarsometatarsal Joint in Relation to Hallux Valgus Deformity: Anatomical and Biomechanical Aspects, *Foot and Ankle International*, Volume: 20 issue: 10, page(s): 651-656.

¹⁵ Luhmann Scott, Rich Margaret, Painful Idiopathic Rigid Flatfoot in Children and Adolescents, *Foot and Ankle International*, Volume: 21 issue: 1, page(s): 59-66, Jan. 2000.

¹⁶ Halabchi Farzin, Mazaheri Reza, Pediatric Flexible Flatfoot; Clinical Aspects and Algorithmic Approach, *Iranian Journal of Pediatrics*, 2013 Jun; 23(3): 247–260.

¹⁷ Lamm Bradley, Stasko Paul, Normal Foot and Ankle Radiographic Angles, Measurements, and Reference Points, *Journal Foot and Ankle Surgery*, Vol. 55, issue 5, 2016.



Dr. Mendeszoon is a foot and ankle surgeon at Precision Orthopaedic Specialties, Inc., a multi-specialty orthopedic practice in Chardon, Ohio. He is the Director of the Advanced Foot and Ankle Surgical Fellowship through University Hospitals—Richmond Medical Center, and Chief of Podiatry at University Hospital's Geauga Medical Center. Dr. Mendeszoon is an assistant clinical professor at the Ohio University Heritage Medical School and an instructor of surgery at

Kent State University School of Podiatric Medicine.

CME EXAMINATION

SEE ANSWER SHEET ON PAGE 143.

1) The human body has approximately how many muscles?

- A) Two hundred
- B) Four hundred
- C) Six hundred
- D) Eight hundred

2) The functional unit of a skeletal muscle is

- A) Sarcomere
- B) Myosin
- C) Endoplasmic reticulum
- D) Myofibril

3) Delayed onset muscle soreness (DOMS) typically occurs how many hours after exercise?

- A) 6 hours
- B) 12 hours
- C) 18 hours
- D) 24 hours

4) Which foot and ankle biomechanical theory is generally NOT utilized in evaluating a patient?

- A) Root theory
- B) Kirby theory
- C) Jones theory
- D) Dananberg theory

5) The majority of overuse conditions develop in which part of the gait cycle?

- A) Swing phase
- B) Heel contact
- C) Midstance
- D) Toe off

6) The human adult walking gait cycle is developed at about the age of _____ ?

- A) One
- B) Two
- C) Four
- D) Six

Continued on page 142

- 7) Over 98% percent of tendons are composed of
- A) Muscle
 - B) Ligaments
 - C) Water
 - D) Type 1 collagen
- 8) _____ deformity is the most common type of foot deformity and is directly related to impaired biomechanics.
- A) Rigid flatfoot
 - B) Flexible flatfoot
 - C) Cavus foot
 - D) Charcot foot
- 9) The etiology of pes plano valgus deformity can be derived from:
- A) Genetics
 - B) Trauma
 - C) Biomechanical factors
 - D) All of the above
- 10) It is recommended that the average person take how many steps a day to remain active and healthy?
- A) 5,000
 - B) 10,000
 - C) 15,000
 - D) 20,000

SEE ANSWER SHEET ON PAGE 143.

The author(s) certify that they have NO affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest), or non-financial interest (such as personal or professional relationships, affiliations, knowledge, or beliefs) in the subject matter or materials discussed in this manuscript.

PM's CME Program

Welcome to the innovative Continuing Education Program brought to you by *Podiatry Management Magazine*. Our journal has been approved as a sponsor of Continuing Medical Education by the Council on Podiatric Medical Education.

Now it's even easier and more convenient to enroll in PM's CE program!

You can now enroll at any time during the year and submit eligible exams at any time during your enrollment period.

CME articles and examination questions from past issues of *Podiatry Management* can be found on the Internet at <http://www.podiatrym.com/cme>. Each lesson is approved for 1.5 hours continuing education contact hours. Please read the testing, grading and payment instructions to decide which method of participation is best for you.

Please call (631) 563-1604 if you have any questions. A personal operator will be happy to assist you.

Each of the 10 lessons will count as 1.5 credits; thus a maximum of 15 CME credits may be earned during any 12-month period. You may select any 10 in a 24-month period.

The Podiatry Management Magazine CME program is approved by the Council on Podiatric Education in all states where credits in instructional media are accepted. This article is approved for 1.5 Continuing Education Contact Hours (or 0.15 CEU's) for each examination successfully completed.

PM's privacy policy can be found at <http://podiatrym.com/privacy.cfm>.

**Home Study CME credits now
accepted in Pennsylvania**

Enrollment/Testing Information and Answer Sheet

Note: If you are mailing your answer sheet, you must complete all info. on the front and back of this page and mail with your credit card information to: **Program Management Services, P.O. Box 490, East Islip, NY 11730.**

TESTING, GRADING AND PAYMENT INSTRUCTIONS

(1) Each participant achieving a passing grade of 70% or higher on any examination will receive an official computer form stating the number of CE credits earned. This form should be safeguarded and may be used as documentation of credits earned.

(2) Participants receiving a failing grade on any exam will be notified and permitted to take one re-examination at no extra cost.

(3) All answers should be recorded on the answer form below. For each question, decide which choice is the best answer, and circle the letter representing your choice.

(4) Complete all other information on the front and back of this page.

(5) Choose one out of the 3 options for testgrading: mail-in, fax, or phone. To select the type of service that best suits your needs, please read the following section, "Test Grading Options".

TEST GRADING OPTIONS

Mail-In Grading

To receive your CME certificate, complete all information and mail with your credit card information to: **Program Management Services, P.O. Box 490, East Islip, NY 11730. PLEASE DO NOT SEND WITH SIGNATURE REQUIRED, AS THESE WILL NOT BE ACCEPTED.**

There is **no charge** for the mail-in service if you have already enrolled in the annual exam CME program, and we receive this exam during your current enrollment period. If you are not enrolled, please send \$27.00 per exam, or \$219 to cover all 10 exams (thus saving \$51 over the cost of 10 individual exam fees).

Facsimile Grading

To receive your CME certificate, complete all information and fax 24 hours a day to 631-532-1964. Your CME certificate will be dated and mailed within 48 hours. This service is available for \$2.50 per exam if you are currently enrolled in the annual 10-exam CME program (and this exam falls within your enrollment period), and can be charged to your Visa, MasterCard, or American Express.

If you are *not* enrolled in the annual 10-exam CME program, the fee is \$27 per exam.

Phone-In Grading

You may also complete your exam by using the toll-free service. Call 1-800-232-4422 from 10 a.m. to 5 p.m. EST, Monday through Friday. Your CME certificate will be dated the same day you call and mailed within 48 hours. There is a \$2.50 charge for this service if you are currently enrolled in the annual 10-exam CME program (and this exam falls within your enrollment period), and this fee can be charged to your Visa, Mastercard, American Express, or Discover. If you are not currently enrolled, the fee is \$27 per exam. When you call, please have ready:

1. Program number (Month and Year)
2. The answers to the test
3. Credit card information

In the event you require additional CME information, please contact PMS, Inc., at **1-631-563-1604.**

ENROLLMENT FORM & ANSWER SHEET

Please print clearly...Certificate will be issued from information below.

Name _____ Email Address _____

Please Print: FIRST MI LAST

Address _____

City _____ State _____ Zip _____

Charge to: Visa MasterCard American Express

Card # _____ Exp. Date _____ Zip for credit card _____

Note: Credit card is the only method of payment. Checks are no longer accepted.

Signature _____ Email Address _____ Daytime Phone _____

State License(s) _____ Is this a new address? Yes No

Check one: I am currently enrolled. (If faxing or phoning in your answer form please note that \$2.50 will be charged to your credit card.)

I am not enrolled. Enclosed is my credit card information. Please charge my credit card \$27.00 for each exam submitted. (plus \$2.50 for each exam if submitting by fax or phone).

I am not enrolled and I wish to enroll for 10 courses at \$219.00 (thus saving me \$51 over the cost of 10 individual exam fees). I understand there will be an additional fee of \$2.50 for any exam I wish to submit via fax or phone.

Over, please

EXAM #3/18
Fundamentals of Care in Biomechanics—Part I
(Mendeszoön)

Circle:

- | | |
|------------|-------------|
| 1. A B C D | 6. A B C D |
| 2. A B C D | 7. A B C D |
| 3. A B C D | 8. A B C D |
| 4. A B C D | 9. A B C D |
| 5. A B C D | 10. A B C D |

This CME activity is free from commercial biases and is under the overall management of *Podiatry Management Magazine*.

Medical Education Lesson Evaluation

Strongly agree [5]	Agree [4]	Neutral [3]	Disagree [2]	Strongly disagree [1]
--------------------------	--------------	----------------	-----------------	-----------------------------

- 1) This CME lesson was helpful to my practice ____
- 2) The educational objectives were accomplished ____
- 3) I will apply the knowledge I learned from this lesson ____
- 4) I will make changes in my practice behavior based on this lesson ____
- 5) This lesson presented quality information with adequate current references ____
- 6) What overall grade would you assign this lesson?
A B C D

How long did it take you to complete this lesson?

_____ hour _____ minutes

What topics would you like to see in future CME lessons?
Please list :
