



# Non-Invasive Diagnosis of Refractory and Chronic Gout

This can be accomplished via various imaging techniques.

BY JOHN COZZARELLI, DEMI TURNER, AND NICHOLAS F. COZZARELLI

## Goals and Objectives

To understand the various modalities utilized in imaging gout in all phases of the disease state, either acute or chronic in nature.

To allow the reader to realize that utilizing various imaging modalities for early intervention leads to better management of gout.

Welcome to Podiatry Management's CME Instructional program. Podiatry Management Magazine is approved by the Council on Podiatric Medical Education as a provider of continuing education in podiatric medicine. Podiatry Management Magazine has approved this activity for a maximum of 1.5 continuing education contact hours. This CME activity is free from commercial bias and is under the overall management of Podiatry Management Magazine.

You may enroll: 1) on a per issue basis (at \$28.00 per topic) or 2) per year, for the special rate of \$229 (you save \$51). You may submit the answer sheet, along with the other information requested, via mail, fax, or phone. You can also take this and other exams on the Internet at [www.podiatrym.com/cme](http://www.podiatrym.com/cme).

If you correctly answer seventy (70%) of the questions correctly, you will receive a certificate attesting to your earned credits. You will also receive a record of any incorrectly answered questions. If you score less than 70%, you can retake the test at no additional cost. A list of states currently honoring CPME approved credits is listed on pg. 136. Other than those entities currently accepting CPME-approved credit, Podiatry Management cannot guarantee that these CME credits will be acceptable by any state licensing agency, hospital, managed care organization or other entity. PM will, however, use its best efforts to ensure the widest acceptance of this program possible.

**This instructional CME program is designed to supplement, NOT replace, existing CME seminars.** The goal of this program is to advance the knowledge of practicing podiatrists. We will endeavor to publish high quality manuscripts by noted authors and researchers. If you have any questions or comments about this program, you can write or call us at: **Program Management Services, P.O. Box 490, East Islip, NY 11730, (631) 563-1604 or e-mail us at [bblock@podiatrym.com](mailto:bblock@podiatrym.com).**

Following this article, an answer sheet and full set of instructions are provided (pg. 136).—Editor

## Introduction

Gout is the most common inflammatory arthropathy worldwide and occurs when sodium urate is converted to monosodium uric acid (MSU) and deposits within the joints. Crystallization in the body occurs when serum uric acid is 6.8mg/dl or higher.<sup>1</sup> In a national health survey conducted in 2007 and 2008,<sup>2</sup> 3 million Americans reported to have a clinical diagnosis of gout.<sup>2</sup> This corresponds to 3.9% of the U.S. adult population. The prevalence of gout in the United States is rapid-

## Gout is the most common mono-arthropathy that occurs worldwide.

ly rising, increasing by 150% in the last decade. Between 2016 and 2017, the number of hospital emissions and emergency visits secondary to uncontrolled gout has doubled.<sup>3</sup>

Gout generally presents as a monoarticular arthritis, though it can be polyarticular and asymmetrical in

distribution. The over-saturation of the MSU crystals in the body interacting with specific immune system cells produces a tophus. Tophi are an abrasive by-product of failed purine metabolism in humans, causing inter-articular and extra-articular damage. While early

*Continued on page 126*

Gout (from page 125)

diagnosis is clinically difficult, it is imperative to prevent permanent and long-term sequelae. The diagnosis of gout has improved due to availability of tests, the ability to definitely identify the MSU crystals in the affected joints via arthrocentesis, and direct fluid examination for crystals.

Lifestyle factors, dietary choices, and socioeconomic parameters have been proposed to explain the more prevalent epidemiological observations. Unhealthy dietary habits and a sedentary lifestyle appear to be major factors for the rise of gout in western countries. Choi, et al. reported a 1.85 relative risk factor for the incidence of gout in males consuming two or more sugar-sweetened soft drinks per day.<sup>4</sup> This is attributed to an increased production of uric acid in the liver, as well as attenuated renal uric acid clearance, both secondary to excessive fructose consumption. Sugar-sweetened soft drinks represent the single largest food source of calories in the U.S. population, and their consumption has consistently increased over the last several decades.<sup>4</sup> This is especially true for fructose and its most popular form HFCS (high fructose corn syrup) that converts to uric acid upon ingestion. Also, this has caused a rise in increased adiposity in the U.S. populace because of the large availability of HFCS in comparison to other forms of fructose.



Figures 1 and 2: Severe chronic gout demonstrating classic sclerotic edges and tophus in soft tissue and "Martel's sign."

With the rise of these co-morbidities, the result has been an increased incidence of gout. Co-morbid conditions including obesity, hypertension, insulin resistance, type 2 diabetes, and

be secondary to the increased consumption of inexpensive sources of animal protein, and concentrated animal protein supplements. There is a larger number of sweeteners available today


**Chronic gout can easily be assessed best by x-ray.**

chronic kidney disease. These are all associated with the metabolic syndrome and are well-established independent risk factors for the occurrence of gout. With the advent of bariatric surgery, these patients are also more prone to gout attacks secondary to increased protein intake with dehydration.<sup>5</sup> The rising prevalence may also

that are metabolically difficult for the kidneys to efficiently process.<sup>4</sup>

This directly yields to a condition of uric acid under excretion in the kidneys. It is thought to be the most relevant failed metabolic action versus the overproduction of serum urate in the body. A 2009-2010 NHANES

*Continued on page 127*



Figures 3, 4, 5, 6: Periarticular, punched out, intra-articular, sclerotic edges.

## Gout (from page 126)

sub-analysis reported a prevalence rate >60% for CKD 3 and CKD 4 in patients with gout. Sustained hyperuricemia may cause renal interstitial urate crystal deposition causing gouty nephropathy. Renal inflammation and renal insufficiency often follow. There is a 22% prevalence of nephrolithiasis among patients with primary gout. Further, 10% of patients had kidney stones (urolithiasis) comprised of urate. Therefore, new techniques for early

detection and treatment of gout will be necessary in managing this rising epidemic, along with pharmacological and lifestyle changes. With advanced imaging techniques, earlier diagnosis can be made, facilitating prompt treatment and intervention, preventing long-term

damage and having the patients achieve clinical remission.<sup>7</sup>

Elevated serum uric acid levels are present in 21% of the U.S. population, but only 5-20% of the individuals with hyperuricemia develop gout. Patients with hyperuricemia may not

## Bone erosion is caused by reduction of osteoblasts and an increase in osteoclast activity.




Figure 7: Sonoscape 16 mg-Hz linear array transducer demonstrating the double contour sign of the first MTPJ in plantar long axis with standoff.



Figure 8: Classic snowstorm effect with standoff in medial long axis position.

necessarily develop gout; and those with gout may have normal serum uric acid levels, particularly during acute gout attacks. Higher degrees and longer duration of elevated urate levels can lead to a greater severity of disease of gouty arthritis, which is the impetus for early and aggressive lowering of urate levels in the treatment of gout to sUA levels below 6 mg/dL. This new “treat to target approach” helps facilitate a positive treatment outcome for a patient.<sup>8</sup>

Urate crystals are deposited predominantly along the surface of the articular cartilage. 65% of the time, gout manifests in the first metatarsal phalangeal joint.<sup>9</sup> 90% of all gout patients will have refractory attacks that occur in the foot. When it occurs in the first metatarsal phalangeal joint, it is known as podagra. Mid-foot (25%-50%) and ankle (18%-60%) attacks can also commonly occur.<sup>1</sup> The total urate burden of the body is comprised of 50% being in the foot and ankle.<sup>23</sup> The current gold standard for the diagnosis of gout is aspiration of the involved joint, followed by polarized light microscopy of the aspirate assessing for needle-shaped negatively birefringent MSU crystals.

Arthrocentesis may technically be challenging in several scenarios, including anatomically difficult-to-access joints such as the spine and sacroiliac joints in certain patients, such as those with immunodeficiencies, patients with inadequate fluid volumes, or patients with severely inflamed joints. In addition, arthrocentesis is a painful invasive technique whose risk factors include infection, hemorrhage, and damage to involved tissues. Experience in diagnosing gout is frequently and definitively made by joint aspiration to rule in the presence of MSU crystals

*Continued on page 128*

*Gout (from page 127)*

as well as being critically important to rule out septic arthritis or an immune modulation response, which may mimic gout clinically. Samples retrieved by arthrocentesis require immediate analysis to preserve the technique's sensitivity and specificity, introducing practical limitations.<sup>10</sup>

The purpose here is to review the imaging appearances of gout in the clinical presentation to assist in diagnosis and to help the clinician understand what he/she is seeing. With the aid of this advanced imaging, the physician can institute a treatment regimen earlier based on achieving a diagnosis earlier than previously had not been available. Many of the diagnostic techniques available today were previously undeveloped or unknown. Imaging may provide a non-invasive and accurate way to diagnose gout. Different imaging modalities may include x-ray, computed tomography (CT), dual energy CT (DECT), magnetic resonance imaging (MRI), and ultrasound (US).

### Incidence and Progression

It is generally recognized that gout prevails in the classical presentation favoring 20:1 men versus women; and in over 80% to 90% of gout patients there is a positive familial history of gout. As women become post-menopausal, the rate of prevalence changes with roughly 7% of men and 5% of women. Post-menopausal women no longer produce estrogen, which is thought to be a uricosuric that suppresses the MSU crystal precipitation response.<sup>11</sup> Also, there is another statistical bump in the women's curve due to the use of thiazide diuretics as a first line medical regimen in hypertension.<sup>12</sup>

Gout the disease is basically divided into four phases:

- 1) Asymptomatic hyperuricemia, 2) Acute gout, 3) Inter-critical gout, and 4) Chronic gout.<sup>13</sup>

Chronic gout may take at least 8 to 10 years to manifest with abnormalities on plain film. Therefore, advanced imaging is frequently employed for early detection, to ascertain a confirmatory diagnosis, and determine the extent of the structural destruction to the joints. By the time the

first acute attack occurs, gout is most likely present for 10 to 12 years.

Sites frequently affected are the first metatarsophalangeal joint in the foot. Other frequent sites are the hand, hind foot, pre-patellar bursa, heart, kidneys and at the insertion points for ligaments and tendons.<sup>14</sup> This frequently can cause severe enthesopathies at these sites and may cause significant disabilities. Some of the common pathological findings of

gout are juxta-articular erosions, cortical depressions, and overhanging edges with sclerotic margins. These findings are seen routinely and often the cortical depressions are adjacent to a tophus. CT is the most sensitive for detecting bony erosions. Secondly, synovial proliferation may be demonstrated on imaging as synovial thickening, often with synovial enhancement.

*Continued on page 129*



Figure 9: Dorsal long axis imaging with standoff with linear array transducer at 16mg-Hz demonstrating synovitis.




Figure 10: Classic tophus identified with sonoscape 16mg-Hz linear array transducer with standoff demonstrating varying degrees of hypo and hyper echogenicity noted in the medial long axis position.

Gout (from page 128)

Contrast-enhanced images may increase detection of bone erosions. US Doppler imaging is utilized in gout to assess for increased blood flow and active inflammation. The tophus is visualized (both intra- and extra-articularly) as a high-density soft tissue mass representing the body's chronic immunological response to the MSU crystals. Interarticular presentation is as a hypochoic return with a halo effect that will have a hyperechoic or heterogeneous center. It is now recognized also in axial skeletal imaging of the spine with micro-tophi being imaged.<sup>15</sup>

The characteristics of cartilaginous deposits are not readily demonstrated with conventional diagnostic imaging

and it can have the accumulation of urate crystal deposition that can often resemble xanthoma.<sup>18</sup> This is found in places such as fingers, ears, or eyelids. Soft tissue findings are present with gouty tophi in about 50% of the population.

Descriptions of bone erosions may have overhanging edges as well as

relatively insensitive to the early manifestations of gout. The sensitivity of radiography is only approximately 30%, compared with arthrocentesis, which is considered a gold standard. Gout is often present in the manner reminiscent of psoriatic arthritis without the usually more advanced demineralization and non-uniform joint space narrowing.

Classic osseous radiographic findings demonstrate well defined "punched out" periarticular erosions with sclerotic overhanging edges, normal mineralization, relative preservation of the joint spaces, and asymmetric poly-articular distribution. Joint space widening and subchondral bone collapse might occur in later stages of the disease. Such find-

and it can have the accumulation of urate crystal deposition that can often resemble xanthoma.<sup>18</sup> This is found in places such as fingers, ears, or eyelids. Soft tissue findings are present with gouty tophi in about 50% of the population.

Descriptions of bone erosions may have overhanging edges as well as

*Continued on page 131*

## The best way to visualize non-visible tophus is via a DECT Scan.

including roentgenography, computed tomography (CT) or magnetic resonance imaging (MRI). These types of imaging modalities can nevertheless provide helpful diagnostic clues. However, disadvantages include lack of specificity (bone scan, MRI), considerable cost (MRI), and inability to assess early soft tissue changes such as joint effusions, early erosions, synovial hypertrophy, and hypervascularity, or small tophi (roentgenography). Typical well-defined, "punched out," periarticular erosions with overhanging edges are generally not seen radiographically until 10-12 years after the initial acute attack.<sup>16</sup> Podagra, gout, and calcium pyrophosphate disorder arthropathy can be evaluated with radiographic and ultrasound imaging for detection and confirmation.


### Radiographic Diagnosis of Gout

Traditionally, plain radiography has been the standard initial imaging technique for assessing structural changes in gouty arthropathy, and a quantitative scoring system has been available since 2007. However, erosive changes on radiographs are a late feature of gout and are frequently only apparent after arthritis has been established for more than 10 years. Radiography is

ings take years to manifest after an acute attack. If these findings are present without urate deposits, then it indicates osseous changes due to remote, currently inactive gout.<sup>17</sup> The presence of tophi with underlying bone erosions may indicate the presence of chronic gout.

In patients with acute gout, there is frequently no radiographic change, apart from non-specific soft tissue swelling in the region of the inflamed joint. The hallmark of cartilage damage is joint space narrowing, and this is typically absent in acute gout, but can occur in patients with OA-associated disease, where adjacent sclerosis and osteophyte formation might also be present.

Radiographic presentation of MSU crystals in the synovial membrane in or around ligaments are present in the hands and feet first at the MTPJ and IPJ. As discussed previously, gout may affect numerous joints throughout the body, including the elbow, wrist, olecranon bursa, shoulder, and hip. Gout can also be present in places such as the ear helix or other fleshy areas of the body,



Figures 1 | A, B, C, D, E:  
 A. Typical hyperechoic enhancement on the cartilage surface ("double contour sign")  
 B. Intra-articular hyperechoic clouds  
 C. Bony erosions  
 D. Synovitis  
 E. Power Doppler assessing vascular status of acute inflammatory effects of gout

**TABLE I:**  
**Comparison of Different Imaging Modalities  
in Patients with Gout**

<b>Imaging Feature</b>	<b>Pain Radiography</b>	<b>Ultrasonography</b>	<b>MRI</b>	<b>CT</b>	<b>DECT</b>
Erosions	Fair: Erosions might not be detected at complex regions (Carpus/tarsus) +	Good: Some regions not accessible ++	Very good: Detection of erosions in all areas ++	Very good: Detection of erosions in all area +++	Very good: Detection of erosions in all area
Synovitis or Tenosynovitis	Not imaged (apart from soft tissue swelling) -	Very good: Especially using PDUS +++	Very good: Single regions only (intravenous gadolinium contrast) +++	Not imaged  ++	Not imaged
Osteitis or Bone marrow edema	Not imaged -	Not imaged -	Very good +++	Not imaged -	Not imaged
Cartilage	Poor: Superficial tophi seen adjacent to erosions as asymmetric masses +	Very good: Tophi can be detected at most sites depending on access +++	Fair: Dependent on sequences and acquisitions  +++	Poor  -	Poor  -
Tophus (detection)	Poor: Superficial tophi seen adjacent to erosions as asymmetric masses +	Very good: Tophi can be detected at most sites depending on access +++	Very good at all sites  +++	Good: Cannot always be distinguished from background tissue ++	Very good: Highly specific; subcutaneous and intraosseous tophi detected
Tophus (vascularity and measurement)	Poor  -	Very good: Vascularity detected using PD US; outlining can measure size and volume +++	Very good: Vascularity detected using Gd-contrast; outlining can measure size +++	Very good: Outlining can measure size (with excellent reproducibility) cannot detect vascularity +++	Very good: Outlining can measure size; cannot detect vascularity
Quantification of urate burden	Poor -	Very good: Site specific only ++	Poor	Poor	Very good +++
Effusion	-	+++	+++	++	++
Synovial Proliferation	-	+++	+++	+	+
Joint Space Narrowing	+++	+++	+++	+++	+++
<b>Overall</b>	Widely used multiple joints imaged; inexpensive; minor radiation exposure	Operator dependent; some joints inaccessible versatile; inexpensive; dynamic and quick no radiation	Single joint area imaged; time consuming; very expensive; claustrophobia; all features well imaged; no radiation	Restricted joint area imaged (but wider than MRI); cannot image inflammation; expensive; radiation exposure	Restricted joint area imaged (but wider than MRI); cannot image inflammation; expensive; radiation exposure; Very specific to urate detection and how to quantify urate burden which is unique to this modality

- Poor    + Fair    ++ Better    +++ Best

# CLINICAL PODIATRY

*Gout (from page 129)*

punched-out rat bite lesions often accompanied by osteitis and synovitis. The tophus is a reaction that contributes to joint arthritis associated with this malady. Early findings of gout are often adjacent to the tophus and present often as an incidental finding; notably, acute gout flares can mimic septic arthritis, which must be ruled out.

The hallmark radiographic sign of gout is Martel's sign. The presence of tophi is associated with increased osteoclastogenesis. In the presence of urate deposition from gout, affected bone will frequently have erosions that have a punched-out appearance with well-defined overhanging edges.

The erosions are due to tophaceous gout of the first metatarsophalangeal joint. It will have typical "overhanging margins" of bone both in the phalanx and in the lateral aspect of the metatarsal head in the case of classic Podagra.<sup>19</sup> The other hallmark sign of gout is the double contour sign imaged via ultrasound (Figures 1 and 2).

### Ultrasonography

Musculoskeletal ultrasound studies are now performed with triple blended frequency linear array transducers (Figures 3-6). Frequencies are between 7-16 MHz. The transducer is 50mm in length. A standoff is typically utilized to conform to irregular surfaces. The standoff pushes the

image down in the near field about 1 cm. This allows better signal to noise and decreases the speckles in the anechoic near field. Standard imaging positioning is distal to the right and proximal to the left. Standard imaging plans are in the short and long axes. All ultrasound exams are operator-dependent and are dynamic in nature. The exams can be reviewed in the cine-loop.

In common with the other advanced modalities, ultrasonography is more sensitive than plain radiography for detecting erosions. Ultrasonography can detect three times as many erosions as radiography, indicating it is superior for sensitivity. There is no

*Continued on page 132*




Figure 12: Dual energy computed tomography (DECT) demonstrated gout, in green, in the hand, feet, elbow, knee, and spine.

*Gout (from page 131)*

gold standard against which to compare ultrasonography.

Power Doppler ultrasonography (PDUS) can detect slow blood flow at the site being imaged, which may indicate inflammatory tissue. Active or “hot” sites will demonstrate PDUS signals. Deposits of MSU crystals can be detected on ultrasonography as tophi and as a fine powder layer over the cartilage, labeled as the “double contour sign”. The double contour sign is defined as a highly echogenic line which is detected parallel to the echogenic line of cortical bone, with the interposed low signal representing cartilage.

Although this sign has not been compared directly with histopathological appearances, it does seem to correlate with findings from a necropsy study where MSU crystal deposits were seen to overlie the cartilage surface in gouty joints. The double contour sign is present roughly 92% of the time in gouty joints. The double contour sign will be present even if

clinical examination. Utilizing the volume calculation feature on ultrasound machines, the tophaceous volume can be calculated to evaluate if urate-lowering therapy is effective. This modality provides the operator the feasibility to discriminatively measure and evaluate

response after the presentation of the acute deposition of the gouty crystals.

Ultrasound can also visualize the insertions of ligaments and tendons for urate crystal deposition in patients with an acute flare or to differentiate other diseases such as osteoarthritis. On ul-




Figure 13 and 14: DECT scan of the feet demonstrating chronic gout (in green). DECT scan of the feet after receiving Krystexxa<sup>®</sup>; gouty tophus is relatively gone.

changes in tophus size.

Other classic patterns observed in the joint effusion of gout are an anechoic cavity. Homogenous punctuate echogenic foci of urate sand can also be observed. The “snowstorm” pattern demonstrates rounded aggregates of variable echogenicity super-

trasound, tophi may demonstrate an anechoic peripheral halo with a heterogeneous hyperechoic center adjacent to a joint and may often be confused with rheumatoid arthritis. Micro-tophi may also be seen in joint effusions, creating hyperechoic foci. This is referred to as a snowstorm appearance. Synovitis and erosions with irregularity of the cortex may be indicative of gout.

Besides detection of synovitis and erosions, ultrasound can also detect the presence of the double contour sign (DC) and peri-intra-articular hyperechoic cloudy areas (HC). DC is defined as focal or diffuse echogenicity on the surface of the joint cartilage. HC is characterized as peri-/intra-articular heterogeneous masses composed of hyper- or hypo-echoic material sometimes possessing posterior acoustic shadows. Synovitis is defined as the presence of either synovial fluid and/or synovial hypertrophy, which is characterized by abnormal hypoechoic intra-articular soft tissue. Volume calculations can be performed both in the long and short axis on a dual screen. The volume calculation occurs with an “x”, “y” and “z” axis measurement.

Area calculations of tophus can be

*Continued on page 133*

### **Martel’s sign is defined as an overhanging C-shaped edge of bone characteristic of gouty arthritis.**

anisotropy is done at the time of imaging.

Although the double contour sign has mostly been described in patients with long-standing disease, it has been reported to occur in a significant proportion of patients with asymptomatic hyperuricemia and no clinical history of musculoskeletal involvement. This sign has been reported to disappear with effective ULT, once sustained normouricemia has been achieved.<sup>20</sup>

Tophi appear on ultrasonography as hyperechoic nodules, often poorly defined contours and surrounded by an anechoic halo (Figure 7). Tophi can be detected subcutaneously or in deep tissues, where they are non-visible on

imposed on urate sand.<sup>21</sup>

Ultrasound presents a rather unique ability to be primed for gout remission management. It can be one of the more sensitive and specific imaging modalities to detect MSU intra-articular crystal deposition and early joint destruction. This is especially due to the ability to utilize ultrasound that is high frequency and high resolution to detect changes in the density of the articular surface and synovial fluid as well as changes in density and thickening of the articular cartilage, giving rise to the double contour sign previously mentioned. It also shows the infiltrate into the soft tissue that causes the itching inflammatory

*Gout (from page 132)*

accomplished by tracing the tophus. Finally, with the aid of different color maps, three-dimensional scanning is accomplished to image the tophus. If a patient is going to be placed on a uricolytic agent such as Krystexxa®, a pre-treatment image and volume calculation of a tophus can be done and post-treatment, the tophus site can be imaged to show resolution of the tophus (Figures 8-13).

## CT

Conventional CT or single energy computed tomography (SECT) is an excellent modality for imaging tophi and erosions in patients with gout, owing to its multi-planar capabilities. It utilizes a single polychromatic x-ray beam ranging from 70 to 140 kVp with a standard of 120 kVp. CT is probably a more accurate tool of detecting and measuring tophi than MRI and ultrasonography because of its superior definition of tophus-bone interface. Tophi have a typical density on CT of 170 Hounsfield units. CT can detect tophi that are not apparent on clinical physical examination. Unfortunately, CT has a low false-positive rate for detecting tophi,

Calcification in the tophus often is suggestive of renal impairment, which indicates that the kidneys are under secretors. While joint effusions may also be seen, they may not be specific for gout unless accompanied by numerous tophi. (Table 1 summarizes these imaging findings).

CT, especially dual energy, can easily distinguish between crystals and calcification, differentiating between biomechanical abnormalities and crystalline deposition. This could be useful when the clinical presentation is unclear or as an atypical presentation for the clinician treating gout.

While CT may be superior to plain film in detecting abnormalities in gout, it is more expensive than x-ray. It provides, however, the information that can be used to determine the aging extent of the pathological manifestation of the erosions of gout. CT images may show the punched out lytic lesions and overhanging sclerotic margins of early bone erosion. CT can allow the clinician to rule out other diseases such as rheumatoid arthritis or CPPD (calcium pyrophosphate disease). In CPPD,

there often is non-uniform joint space narrowing, subchondral sclerosis of the joint, and peri-articular soft tissue swelling due to tophi being deposited around the joints.

## DECT

The newest modality in imaging gout is dual-energy computed tomography (DECT). It is also known as spectral imaging. An image is acquired simultaneously using two ray tubes.

Two different voltages are utilized, such as 80 kVp and 140 kVp. The beams are 90 degrees to each other. Utilizing this technique, different materials such as bone, soft tissue, and gouty tophus may be discriminated based on their varying absorption of different energies depending on their chemical composition. (22)

DECT is very sensitive and specific for its identification of tophi. However,

*Continued on page 134*




Figure 15: Short axis and long axis scan demonstrating volume of tophus.




Figure 16: 3D image demonstrating tophus of the first MTP] clearly demonstrating bone erosions by tophus. Image can be independently rotated on the x,y,and z axis.

Gout (from page 133)

utilization of DECT imaging may be limited due to the cost and radiation. DECT has the ability to show intra- and extra-articular tophi as well as tophi located at the insertions of the ligaments and tendons, and within the spine. DECT scanning allows the physician to ascertain whether the patient is tophaceous, even if it is not visible on examination.<sup>23</sup> This technique utilizes ionizing radiation and is relatively expensive, so it might not be widely applicable outside of research or teaching hospitals. With this technique, there is a 10% increase in radiation exposure compared to a standard CT examination (Figure 14).

### MRI

Magnetic resonance imaging may also be used in the diagnosis of gout (Figures 15 and 16). Most tophi are heterogenous in signal secondary to their varying high-protein composition. MRI allows accurate crystalline deposit identification and localization, including within the deeper soft tissues and in areas and in sites atypical for MSU and/or CPPD crystalline deposition. It can allow the clinician to diagnose the extent of gout involvement, including at the insertions of tendons and ligaments into bone.

Gouty tophi, when examined with T1-weighted imaging post-contrast, frequently yield an image of good quality to clearly demarcate the size borders and depth of penetration of the tophus.

A T2-weighted imaging can have a higher signal to noise ratio post-contrast. Detected signal strength is in the mean proportional to the intensity of bone involvement in the structure, especially in the inter-osseous layers of the foot due to the sensitivity of MRI.

MRI modality can utilize fat suppression imaging. With MRI, the progress of treatment for chronic refractory gout can be monitored and assessed continuously. Additionally, with MRI, gout and other crystalline arthropathies such as CPPD can be frequently differentiated between septic arthritis and osteoarthritic conditions of the joint (Figure 17).<sup>24</sup>

### Conclusion

Gout is an erosive inflammatory arthropathy where long-term damage to joints can have a major negative impact on patient function. Cellular studies have focused on the role of cytokines and inflammatory mediators produced by cells of the innate and adaptive immune systems that are resident within the tophi. MSU crystals can act both directly and indirectly to stimulate an inflammatory response with resultant tissue destruction.

Osteoclasts and osteoblasts in the underlying bone have recently




Figure 17: MRI of the foot demonstrating gouty tophus with erosions of the first MTPJ and phalanx.

emerged as major players in driving erosion and preventing adequate tissue repair when subjected to an environment of MSU crystal stimulation. Advanced imaging modalities provide additional insights into the pathological processes involved in tissue destruction in patients with gout by providing information about bone erosions, tophi, and MSU deposition on cartilage surfaces. All modalities have advantages and disadvantages. Thus, insights from cellular and imaging studies can complement one another and help improve the clinical management of patients with this painful and potentially disabling disease (Figure 18). **PM**

### References

- Roddy, E. *J Foot Ankle Res.* 2011;4:1-6.
- Krishnan E, et al. *J Rheumatol.* 2008;35(3):498-501.
- Lim SY, et al. *JAMA.* 2016;315:2345-2347.
- Choi HK, Curhan G. Soft Drinks, fructose consumption, and the risk of gout in men: prospective study. *BMJ.* 2008;336(7639):309-312.
- Tana, Claudio C. "Management of Hyperuricemia and Gout in Obese Patients Undergoing Bariatric Surgery." *Postgraduate Medicine*, vol. 130, no. 6, 22 June 2018, pp. 523-535., doi:10.1080/00325481.2018.1485444.
- "Reduced glomerular function and prevalence of gout: NHANES 2009-10" *PLoS one* vol. 7,11 (2012): e50046.
- Parikh P, et al. *J Rheumatol.* 2010; 37:2190-2191.
- Perez-Ruiz F, Lioté F. Lowering serum uric acid levels: what is the optimal target for improving clinical outcomes in gout? *Arthritis Rheum.* 2007;57(7):1324-1328.
- Jelley, M J MJ. "Practical Steps in the Diagnosis and Management of Gout." *BioDrugs*, vol. 14, no. 2, Aug. 2000, pp. 99-107., doi:14(2):99-107.
- Taylor WJ, Fransen J, Dalbeth N, et al.

Continued on page 135




Figure 18: Montage showing MRI, x-ray, ultrasound, and DECT scanning.

## Gout (from page 134)

Diagnostic arthrocentesis for suspicion of gout is safe and well tolerated. *J Rheumatol.* 2016;43(1):150-153.

<sup>11</sup> Bruderer, Saskia G SG. "Association of Hormone Therapy and Incident Gout: Population-Based Case-Control Study." *Menopause*, vol. 22, no. 12, Dec. 2015, pp. 1335-42., doi:10.1097.

<sup>12</sup> Ranieri, Laura et al. "Impact of diuretics on the urate lowering therapy in patients with gout: analysis of an inception cohort" *Arthritis research & therapy* vol. 20, 1 53. 22 Mar. 2018, doi:10.1186/s13075-018-1559-2.

<sup>13</sup> Brook RA, Forsythe A, Smeeding JE, Lawrence EN. Chronic gout: epidemiology, disease progression, treatment and disease burden. *Curr Med Res Opin.* 2010;26:2813-21.

<sup>14</sup> Edwards, N. Lawrence., Terkeltaub, Robert. "Gout: Diagnosis and Management of Gouty Arthritis and Hyperuricemia." *Gout: Diagnosis and Management of Gouty Arthritis and Hyperuricemia, Professional Communication*, 2016, p. 91.

<sup>15</sup> Konatalapalli RM, Demarco PJ, Jelinek JS, et al. Gout in the axial skeleton. *J Rheumatol.* 2009;36:609-613.

<sup>16</sup> Haseeb A, et al. *Turk Rheumatol.* 2012;27:208-211.

<sup>17</sup> RheumTutor.com. <http://www.rheumtutor.com/xray-findings-in-gout-2/>. Accessed October 11, 2018.

<sup>18</sup> Sutton, Leigh L. "Perforating Gout of the Ear." *Dermatology Online Journal*, vol. 22, no. 10, 15 Oct. 2016, pp. 1087-2108.

<sup>19</sup> Martel W. *Radiology.* 1968;91:755-756.

<sup>20</sup> Thiele, R. G., and N. Schlesinger. "Diagnosis of Gout by Ultrasound." *Rheumatology*, vol. 46, no. 7, July 2007, pp. 1116-1121.

<sup>21</sup> Grassi W, et al. *Semin Arthritis Rheum.* 2006;36(3):197-202.

<sup>22</sup> Baer An, Kurano T, Thakur UJ, et al. Dual-energy computed tomography has limited sensitivity for non-tophaceous gout: a comparison study with tophaceous gout. *BMC Musculoskel Discord.* 2016;17(1):91.

<sup>23</sup> Choi HK, et al. *Ann Rheum Dis.* 2009;68:1609-1612.

<sup>24</sup> Chhana, Ashika A. "Advanced Imaging Assessment of Gout: Comparison of Dual-Energy CT and MRI with Anatomical Pathology." *Annals of the Rheumatic Diseases*, vol. 77, no. 4, 2018, pp. 629-630.

**Dr. John Cozzarelli** is in private practice in Belleville, NJ. He is a past adjunct professor in Radiology at NYCPM and is co-founder of the Gout Institute of America.

**Dr. Turner** is in private practice in Montclair, NJ. He is a past associate professor in radiology at NYCPM and is co-founder of the Gout Institute of America.

**Nicholas Francis Cozzarelli** is a 4th year pre-med student at Seton Hall University.



## CME EXAMINATION

SEE ANSWER SHEET ON PAGE 137.

1) Chronic gout can easily be assessed best by:

- A) X-ray
- B) CT scan
- C) DECT scan
- D) Ultrasonography

2) Bone erosion is caused by:

- A) Homeostasis between osteoblasts and osteoclasts.
- B) Reduction of osteoblasts and an increase in osteoclast activity.
- C) Macrophages.
- D) Neutrophils.

3) DECT Scans utilize dual energy sources at:

- A) 45-55 kVp.
- B) 25-55 kVp.
- C) 80-140 kVp.
- D) 75-100 kVp.

4) The best way to visualize non-visible tophus is via a(n):

- A) X-ray.
- B) Ultrasonography.
- C) DECT Scan.
- D) CT scan.

5) The most common mono-arthropathy that occurs worldwide is:

- A) Pseudogout
- B) Psoriatic arthritis
- C) Gout
- D) Rheumatoid arthritis

6) The latest approach to managing gout is described as:

- A) Treating the symptom.
- B) Maintaining serum uric acid above 6.8mg/dl.
- C) Treating to target.
- D) Only using x-ray for diagnosis.

Continued on page 136

- 7) Martel's sign is defined as:
- A) Osteoporotic lesions diagnosed by a DEXA scan.
  - B) An overhanging C-shaped edge of bone characteristic of gouty arthritis.
  - C) Characteristic of refractory gout.
  - D) Occurring in the pelvis.
- 8) When gout occurs in the foot it is called:
- A) King's sign.
  - B) Podagra.
  - C) Locusta sign.
  - D) Carnis sign.
- 9) 90% of gout is caused by:
- A) Overproduction of serum uric acid.
  - B) Under-excretion of uric acid by the respiratory system.
  - C) Under-excretion by the kidneys.
  - D) Subluxation of the spine.
- 10) The only technique that can demonstrate birefringence is:
- A) X-ray
  - B) DECT scan
  - C) Joint aspiration
  - D) Fluoroscopic guidance

**SEE ANSWER SHEET ON PAGE 137.**

The author(s) certify that they have NO affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest), or non-financial interest (such as personal or professional relationships, affiliations, knowledge, or beliefs) in the subject matter or materials discussed in this manuscript.

# PM's CME Program

Welcome to the innovative Continuing Education Program brought to you by *Podiatry Management Magazine*. Our journal has been approved as a sponsor of Continuing Medical Education by the Council on Podiatric Medical Education.

**Now it's even easier and more convenient to enroll in PM's CE program!**

You can now enroll at any time during the year and submit eligible exams at any time during your enrollment period.

**CME articles and examination questions from past issues of *Podiatry Management* can be found on the Internet at <http://www.podiatrym.com/cme>.** Each lesson is approved for 1.5 hours continuing education contact hours. Please read the testing, grading and payment instructions to decide which method of participation is best for you.

Please call (631) 563-1604 if you have any questions. A personal operator will be happy to assist you.

Each of the 10 lessons will count as 1.5 credits; thus a maximum of 15 CME credits may be earned during any 12-month period. You may select any 10 in a 24-month period.

***The Podiatry Management Magazine CME program is approved by the Council on Podiatric Education in all states where credits in instructional media are accepted. This article is approved for 1.5 Continuing Education Contact Hours (or 0.15 CEU's) for each examination successfully completed.***

PM's privacy policy can be found at <http://podiatrym.com/privacy.cfm>.

This CME is valid for CPME-approved credits for three (3) years from the date of publication.

# Enrollment/Testing Information and Answer Sheet

**Note:** If you are mailing your answer sheet, you must complete all info. on the front and back of this page and mail with your credit card information to: **Program Management Services, P.O. Box 490, East Islip, NY 11730.**

## TESTING, GRADING AND PAYMENT INSTRUCTIONS

(1) Each participant achieving a passing grade of 70% or higher on any examination will receive an official computer form stating the number of CE credits earned. This form should be safeguarded and may be used as documentation of credits earned.

(2) Participants receiving a failing grade on any exam will be notified and permitted to take one re-examination at no extra cost.

(3) All answers should be recorded on the answer form below. For each question, decide which choice is the best answer, and circle the letter representing your choice.

(4) Complete all other information on the front and back of this page.

(5) Choose one out of the 3 options for testgrading: mail-in, fax, or phone. To select the type of service that best suits your needs, please read the following section, "Test Grading Options".

## TEST GRADING OPTIONS

### Mail-In Grading

To receive your CME certificate, complete all information and mail with your credit card information to: **Program Management Services, P.O. Box 490, East Islip, NY 11730. PLEASE DO NOT SEND WITH SIGNATURE REQUIRED, AS THESE WILL NOT BE ACCEPTED.**

There is **no charge** for the mail-in service if you have already enrolled in the annual exam CME program, and we receive this exam during your current enrollment period. If you are not enrolled, please send \$28.00 per exam, or \$229 to cover all 10 exams (thus saving \$51 over the cost of 10 individual exam fees).

### Facsimile Grading

To receive your CME certificate, complete all information and fax 24 hours a day to 1631-532-1964. Your CME certificate will be dated and mailed within 48 hours. This service is available for \$2.95 per exam if you are currently enrolled in the annual 10-exam CME program (and this exam falls within your enrollment period), and can be charged to your Visa, MasterCard, or American Express.

If you are *not* enrolled in the annual 10-exam CME program, the fee is \$28 per exam.

### Phone-In Grading

You may also complete your exam by using the toll-free service. Call 1-800-232-4422 from 10 a.m. to 5 p.m. EST, Monday through Friday. Your CME certificate will be dated the same day you call and mailed within 48 hours. There is a \$2.95 charge for this service if you are currently enrolled in the annual 10-exam CME program (and this exam falls within your enrollment period), and this fee can be charged to your Visa, Mastercard, American Express, or Discover. If you are not currently enrolled, the fee is \$28 per exam. When you call, please have ready:

1. Program number (Month and Year)
2. The answers to the test
3. Credit card information

In the event you require additional CME information, please contact PMS, Inc., at **1-631-563-1604.**

## ENROLLMENT FORM & ANSWER SHEET

Please print clearly...Certificate will be issued from information below.

Name \_\_\_\_\_ Email Address \_\_\_\_\_

Please Print: FIRST MI LAST

Address \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Charge to:  Visa  MasterCard  American Express

Card # \_\_\_\_\_ Exp. Date \_\_\_\_\_ Zip for credit card \_\_\_\_\_

**Note: Credit card is the only method of payment. Checks are no longer accepted.**

Signature \_\_\_\_\_ Email Address \_\_\_\_\_ Daytime Phone \_\_\_\_\_

State License(s) \_\_\_\_\_ Is this a new address? Yes  No

**Check one:**  I am currently enrolled. (If faxing or phoning in your answer form please note that \$2.95 will be charged to your credit card.)

I am not enrolled. Enclosed is my credit card information. Please charge my credit card \$28.00 for each exam submitted. (plus \$2.95 for each exam if submitting by fax or phone).

I am not enrolled and I wish to enroll for 10 courses at \$229.00 (thus saving me \$51 over the cost of 10 individual exam fees). I understand there will be an additional fee of \$2.95 for any exam I wish to submit via fax or phone.

Over, please



**EXAM #3/19**  
**Non-Invasive Diagnosis of Refractory and Chronic Gout (J. Cozzarelli, Turner, N. Cozzarelli)**

**Circle:**

- |            |             |
|------------|-------------|
| 1. A B C D | 6. A B C D  |
| 2. A B C D | 7. A B C D  |
| 3. A B C D | 8. A B C D  |
| 4. A B C D | 9. A B C D  |
| 5. A B C D | 10. A B C D |

**Medical Education Lesson Evaluation**

Strongly agree [5]	Agree [4]	Neutral [3]	Disagree [2]	Strongly disagree [1]
--------------------------	--------------	----------------	-----------------	-----------------------------

- 1) This CME lesson was helpful to my practice \_\_\_\_
- 2) The educational objectives were accomplished \_\_\_\_
- 3) I will apply the knowledge I learned from this lesson \_\_\_\_
- 4) I will make changes in my practice behavior based on this lesson \_\_\_\_
- 5) This lesson presented quality information with adequate current references \_\_\_\_
- 6) What overall grade would you assign this lesson?  
A B C D
- 7) This activity was balanced and free of commercial bias.  
Yes \_\_\_\_ No \_\_\_\_
- 8) What overall grade would you assign to the overall management of this activity?  
A B C D

How long did it take you to complete this lesson?

\_\_\_\_ hour \_\_\_\_ minutes

What topics would you like to see in future CME lessons?  
Please list :

---

---

---

---

---