

# Radiology of Systemic Disorders Affecting the Foot and Ankle

Here's a review of the findings seen on foot and ankle x-rays.

BY ROBERT A. CHRISTMAN, DPM, EDM

## Goals and Objectives

After completing this CME, the reader should be able to:

- Discuss and recognize the radiographic features of osteopenia.
- Distinguish between the radiographic findings associated with chronic and regional osteoporosis.
- Describe and identify the radiographic features of metabolic, endocrine, and nutritional disorders that affect the lower extremity.
- Describe and recognize the characteristic radiographic appearance of skeletal dysplasias seen in the lower extremity.
- List disorders that are associated with metastatic calcification, calcinosis, and dystrophic calcification.

Welcome to Podiatry Management's CME Instructional program. Podiatry Management Magazine is approved by the Council on Podiatric Medical Education as a provider of continuing education in podiatric medicine. Podiatry Management Magazine has approved this activity for a maximum of 1.5 continuing education contact hours. This CME activity is free from commercial bias and is under the overall management of Podiatry Management Magazine.

You may enroll: 1) on a per issue basis (at \$35.00 per topic) or 2) per year, for the special rate of \$299 (you save \$51). You may submit the answer sheet, along with the other information requested, via mail, fax, or phone. You can also take this and other exams on the Internet at [podiatrym.com/cme](http://podiatrym.com/cme).

If you correctly answer seventy (70%) of the questions correctly, you will receive a certificate attesting to your earned credits. You will also receive a record of any incorrectly answered questions. If you score less than 70%, you can retake the test at no additional cost. Other than those entities currently accepting CPME-approved credit, Podiatry Management cannot guarantee that these CME credits will be acceptable by any state licensing agency, hospital, managed care organization or other entity. PM will, however, use its best efforts to ensure the widest acceptance of this program possible.

**This instructional CME program is designed to supplement, NOT replace, existing CME seminars.** The goal of this program is to advance the knowledge of practicing podiatrists. We will endeavor to publish high quality manuscripts by noted authors and researchers. If you have any questions or comments about this program, you can e-mail us at [bblock@podiatrym.com](mailto:bblock@podiatrym.com).

Following this article, an answer sheet and full set of instructions are provided.—**Editor**

**M**any systemic disorders demonstrate associated radiographic findings in the lower extremity. Although these conditions typically are not clinically diagnosed by their foot or ankle manifestations, the practitioner should be familiar with these radiologic presentations so they are not misdiagnosed as other conditions. The metabolic diseases, in particular osteoporosis,

include several endocrine and nutritional disorders (hyperparathyroidism, acromegaly, scurvy, and rickets, for example). Paget's disease may be localized in the lower extremity. Skeletal dysplasias present characteristic presentations that may be recognized as incidental findings; examples include osteopetrosis, osteopoikilosis, osteochondromatosis, and osteogenesis imperfecta. Finally, soft tissue manifestations of systemic disorders are

frequently encountered in the foot and leg.

Decreased bone density, or increased radiolucency of bone,<sup>1,2</sup> is a prominent radiographic feature of numerous metabolic disorders. Misconceptions abound regarding the use of terms relating to this finding;<sup>3</sup> therefore, the following definitions are strictly adhered to in this article. The term osteopenia is used to refer to the non-specific radiographic finding of decreased bone

density. The terms decalcification, under-mineralization, and demineralization are not used, because they refer more specifically to the underlying physiology and pathology of bone.<sup>4</sup> Moreover, the term osteoporosis is reserved for the clinical entity.

Generalized, diffuse osteopenia is a highly subjective radiographic finding; one must be careful with this “finding” because radiographic technique and/or processing can profoundly influence the visual appearance (that is, the darkness or lucency) of bones. For example, a dark film can result from high-kVp technique, high-mAs technique. For those still using film and chemical processing, increased time in the developer, or raised developer temperature can result in a dark image. Instead, observe specific, more objective features involving the cortices and/or cancellous bone of the second, third, and fourth metatarsals in the dorsoplantar (DP) foot view. Patterns associated with cortical bone thinning in generalized (chronic, long-standing) osteopenia include endosteal resorption, intracortical tunneling, and/or subperiosteal resorption<sup>1</sup> (Figure 1).



Figure 1: Chronic osteopenia, cortical bone. A, Normally, the endosteal and subperiosteal surfaces of the cortex are well defined and continuous and the remainder of the cortex is radiopaque and homogeneous in density. B, In chronic cortical osteopenia, the endosteal surfaces are ill defined, and lucent striations (intracortical tunneling) may be seen running through the cortex, parallel to the shaft. C, Endosteal resorption, resulting in cortical thinning.



Figure 2: Chronic osteopenia, cancellous bone. A, Cancellous bone is made of primary and secondary trabeculae. The primary trabeculae—also known as stress trabeculae—are found along lines of stress. Secondary trabeculae are found perpendicular or oblique to the primary or stress trabeculae. Secondary trabeculae give spongiosa a fine, homogeneous appearance. B, With chronic osteopenia, secondary trabeculae are resorbed, leaving the primary trabeculae to stand out in relief. As bone is laid down on the remaining primary trabeculae, they become coarse in appearance.



Figure 4: Regional osteoporosis. A, Periarticular, spotty (acute) osteopenia of cancellous bone at metatarsophalangeal joints. B, Metaphyseal ill-defined transverse bands of decreased density, all metatarsals (acute osteopenia). C, Severe acute osteopenia, with spotty, permeative decreased density of cancellous and cortical bone, and subperiosteal resorption. D, Severe osteopenia secondary to polio. Findings parallel those seen in chronic osteopenia (generalized osteoporosis).

A cancellous bone pattern associated with generalized osteopenia is prominent primary trabeculations, which may stand out in relief because secondary trabeculae are resorbed and subsequent bone is laid down on the remaining primary trabeculae (Figure 2). Cortical and cancellous bone resorption is best identified using the DP foot view and concentrating on the second, third, and fourth metatarsals. The medial oblique view should not be used; striations that mimic intracortical tunneling normally are seen at the diaphyseal region. Caution should be exercised so that these findings are not used to diagnose osteoporosis or other metabolic disease. Chronic osteopenia is commonly associated with systemic disorders, which are primarily diagnosed by other, more characteristic, radiologic features.



Figure 3: Generalized osteoporosis. This elderly, postmenopausal patient demonstrates the characteristic features of chronic osteopenia, including endosteal resorption, intracortical tunneling, and prominent, coarsened primary trabeculae.

In contrast, the primary radiographic feature of acute osteopenia is spotty (mottled, moth-eaten) loss of bone density, particularly in periarticular regions. Acute osteopenia is associated with causes of regional osteoporosis (discussed next), such as immobilization or disuse.

**Osteoporosis**

Osteoporosis is a metabolic disease that accompanies many other disease processes, including most of the endocrine and nutritional disorders discussed later in this article. It is characterized by progressive loss of bone mass and may result in pathologic fracture. Osteoporosis is not easily diagnosed with plain films, because 30% to 50% of bone calcium must be lost before osteopenia is visually apparent. Quantitative methods are necessary to detect early osteoporosis; studies include single-energy x-ray absorptiometry (wrist and heel), radiographic absorptiometry (hand), single-photon absorptiometry (wrist), quantitative computed tomography (spine), and dual-photon absorptiometry (spine, hip, body). It has been suggested that the appearance of calcaneal trabecular patterns provide an index for assessing

osteoporosis;<sup>5</sup> however, controversy exists as to whether or not these trabecular patterns correlate with actual bone density.<sup>6</sup> Osteoporosis has been classified according to its etiology: generalized, regional, and localized.<sup>1</sup>

Generalized osteoporosis affects the entire skeleton and is associated with aging (senile osteoporosis), post-menopause, medications (steroids), endocrine states (hyperparathyroidism, diabetes mellitus), deficiency states (scurvy, malnutrition), anemia, and alcoholism, to name a few. The radiographic features are more commonly visualized in the axial skeleton. Typical radiographic findings in the lower extremity (Figure 3) include one or any combination of the following:

- Prominent primary trabeculations
- Thinning of the cortices
- Intracortical striations (tunneling)

Normal radiographic anatomy may be misinterpreted as osteopenia. For example, the trabeculations in the first-metatarsal head and neck are normally prominent and appear coarsened.

Regional osteoporosis affects one extremity. It is associated with disuse (immobilization) and Complex Regional Pain Syndrome (formerly known as reflex sympathetic dystrophy (RSD)). Typical radiographic findings (Figure 4) include one or any combination of the following:

- Spotty (moth-eaten, mottled, patchy) osteopenia in cancellous bone, especially periarticular regions
- Ill-defined transverse bands of decreased density at subchondral or metaphyseal locations

Subperiosteal bone resorption (in severe cases, especially CSDS/RSD)

Long-standing disuse (paralysis is often associated with coarse, prominent primary trabeculations and cortical thinning)

**Normal radiographic anatomy may be misinterpreted as osteopenia.**




Figure 5: Rickets. A, Knee, lateral view. Characteristic features include an ill-defined, lucent and frayed zone of provisional calcification, with widening of the physis. B, Knee, anteroposterior (AP) view. Following treatment, the zone of provisional calcification has become more defined, although still irregular and ill-defined medially. Rickets. C, Ankle, lateral view. Same date as A; similar findings demonstrated, with additional metaphyseal cupping. D, Ankle, lateral view. Metaphyseal cupping very prominent following treatment; also note defect along surface of talar dome.



Figure 6: Hyperparathyroidism. Subperiosteal bone resorption along the medial aspects of the metatarsal shafts.

Figure 7: Renal osteodystrophy. Soft tissue calcification throughout the leg.



Figure 8: Acromegaly. Characteristic features include enlargement of bones (especially metatarsal heads and shafts) and soft tissue thickening (increased soft tissue density and volume).




Figure 9: Scurvy. A, Ankle. Note the transverse line of increased density along the metaphysis. B, Knee (different patient). In addition to the sclerotic line across the metaphysis, beaks are visible at the metaphyseal margins, and Wimberg's sign is evident.

Localized osteoporosis affects one bone. Examples of pathology demonstrating localized osteoporosis include osteomyelitis, arthritis, and neoplasm. The medial aspect of the fifth-metatarsal head is normally radiolucent and should not be mistaken as pathology.

### Osteomalacia and Rickets

Osteomalacia and rickets are characterized histologically as disorders with excessive amounts of uncalcified osteoid.<sup>7</sup> Etiologies include vitamin D deficiency and hypophosphatemia. When present in the adult skeleton, osteomalacia is the name given to this condition; in the growing skeleton of infancy and childhood, it is called rickets.

The radiographic features of osteomalacia are non-specific, consisting primarily of osteopenia. Bowing deformity of long tubular bones may occur. Transversely ori-



Figure 10: Paget's disease. A, Mottise view; B, lateral view. The bone is predominantly sclerotic in the diaphysis and metaphysis, with mixed well-defined lucent areas intermixed. Probably the third stage. (Accompanying text on following page.)

ented, incomplete radiolucencies, known as pseudofractures or Looser's zones, may be seen in tubular bones.<sup>1</sup>

The radiographic features of rickets consist of both non-specific and characteristic findings (Figure 5). Non-specific findings include general retardation of body growth, osteopenia, and bowing deformity of long tubular bones. Characteristic changes occur

## The characteristic feature of hyperparathyroidism is subperiosteal bone resorption.

at the growth plate region of tubular bones and include widening of the physis, decreased density at the zone of provisional calcification, irregularity of the physeal margin of the zone of provisional calcification (which has been described as "fraying" and a "paint-brush appearance"), and widening and cupping of the metaphysis.<sup>1</sup>

### Hyperparathyroidism

The general term hyperparathyroidism refers to increased levels of parathyroid hormone.<sup>1</sup> The characteristic feature of hyperparathyroidism is subperiosteal bone resorption (Figure 6). Other sites of bone resorption include periarticular, intracortical, endosteal, subchondral, and at entheses.

Soft tissue calcification and geographic, lucent lesions known as Brown tumors may also be seen.

### Renal Osteodystrophy

Patients with chronic renal failure demonstrate bony abnormalities known as renal osteodystrophy.<sup>1</sup> The radiographic findings manifested may parallel those of hyperparathyroidism, osteoporosis, and osteomalacia (or rickets if in the child). Calcification of soft tissue and vessels is a frequent finding (Figure 7).

### Acromegaly

Excess pituitary somatotrophic growth hormone results in abnormal growth of bone, cartilage, and fibrous



Figure 11: Hereditary multiple exostoses. A, DP view, left foot. Small exostoses are becoming visible along the medial and lateral aspects of the first-metatarsal distal metadiaphysis and along the lateral aspects of the third- and fourth-toe proximal phalangeal bases. B, Same patient, DP view, right foot. Significant involvement of the second and third metatarsals, with shortening of the second. Hereditary multiple exostoses. C, Same patient, AP view, ankle. Exostoses are seen along the medial aspect of the distal fibular diaphysis and the medial and lateral aspects of the tibial diaphysis. D, Different patient, adult. Involvement of the second through fourth metatarsals and third-toe proximal phalanx.



Figure 12: Osteogenesis imperfecta. A, DP view, foot. Characteristic features include narrowed girth of tubular bones and cancellous osteopenia. B-D, Different patient demonstrating severe osteoporosis and severe deformity of the fibula. Osteogenesis imperfecta. A, DP view, foot. Characteristic features include narrowed girth of tubular bones and cancellous osteopenia. B-D, Different patient demonstrating severe osteoporosis and severe deformity of the fibula.

tissue.<sup>7</sup> In the adult, this is known as acromegaly; gigantism results in the child with open growth plates. Radiographically (Figure 8), the patient with acromegaly has an increase in skin volume; the heel pad thickness is greater than 25 mm in the male and 23 mm in the female (when local causes are excluded). The joint spaces are widened by cartilage thickening. Bones become prominent; the metatarsal heads and distal phalanx ungual tuberosities are enlarged, metatarsal shafts are thickened, and spurs are found at entheses. Interestingly, however, the proximal phalangeal shafts appear narrow in girth.



Figure 13: Osteopetrosis. A, DP view, 7-year-old; B, Lateral view, same patient at 3 years of age. Classic bone-within-bone appearance.



**Hypovitaminosis C (Scurvy)**

Scurvy is caused by insufficient dietary intake of vitamin C.<sup>8</sup> The characteristic radiographic features (Figure 9) occur in the developing child. In the metaphyseal region they include a transverse line of increased density, a transverse line of de-

cessive and abnormal remodeling of bone.<sup>1</sup> Its etiology is unknown, and occurs more frequently in people who are 40 years and older. Paget's disease is typically asymptomatic in the foot, which may explain the infrequent reports at this location in the literature.<sup>9</sup> Four radiographic stages have been recognized. The first stage is described as osteolytic. It starts in the subchondral area, spreading to the metaphysis and eventually the diaphysis resulting in what has been described as a "blade of grass" appearance. The second stage demonstrates both osteolysis and osteosclerosis. The diaphysis appears lucent, whereas the epiphyseal and metaphyseal regions are sclerotic. Bone in the third stage is predominantly sclerotic. Findings include cortical thickening, bone enlargement, and coarsened trabeculae. The fourth phase is malignant degeneration, reported in 1% to 10% of patients.<sup>8</sup> Other complications associated with Paget's disease are fracture, osteomyelitis, and joint disease.<sup>1</sup>

**Hereditary Multiple Exostoses (Osteochondromatosis)**

Hereditary multiple exostoses is



Figure 14: Melorheostosis. The wavy, thickened endosteal cortical surface simulates wax flowing down a candle. This patient also has osteopoikilosis, as demonstrated by the circular increased densities in periarticular regions.



Figure 15: Osteopoikilosis. Multiple islands of cortical bone density are identified in the lesser-metatarsal heads.



Figure 16: Osteopathia striata. Lateral view, ankle. Mixed linear bands of increased and decreased density run through the meta-diaphyseal region of the distal tibia, parallel to the bone's long axis.




Figure 17: Fibrous dysplasia. The geographic lesion in the distal fibular diaphysis demonstrates a hazy, ground glass appearance centrally. Notice its intramedullary location.



Figure 18: Metastatic calcification: secondary hyperparathyroidism. The spectrum of radiographic abnormality may be seen in these patients. This example demonstrates significant soft tissue calcification in the distal leg.

**The radiographic finding described as a "blade of grass" appearance is associated with Paget's disease.**

creased density adjacent to this line of increased density (the scurvy line), and a small, beaklike outgrowth of bone along the margins of the metaphysis. Extensive periostitis may be seen along the entire length of the bone. The epiphysis appears as an outer shell of increased density surrounding a central lucency (Wimberger's sign).<sup>1</sup>

**Paget's Disease (Osteitis Deformans)**

Paget's disease (Figure 10) is characterized by ex-

a skeletal dysplasia resulting from a disturbance of chondroid production resulting in heterotopic proliferation of epiphyseal chondroblasts.<sup>10</sup> The individual lesions appear identical to the bone tumor osteochondroma, that is, multiple cartilage-capped exostoses adjacent to the diaphyseal side of the physis (Figure 11). This condition is autosomal dominant and appears in the first two decades of life as painless bumps near the ends of long bones. Its distribution is frequently symmetric, and it can cause bone deformity and shortening. Approximately 5 percent of these lesions transform into chondrosarcoma.<sup>7</sup>

### Osteogenesis Imperfecta

An inherited skeletal dysplasia that is the result of abnormal metaphyseal and periosteal ossification caused by deficient osteoid production is osteogenesis imperfecta.<sup>10</sup> Abnormal maturation of collagen occurs in both mineralized and non-mineralized tissues. Radiographic features include diffuse osteopenia (occasionally coarse trabeculae and a honeycomb appearance), diminished bone girth, and flared metaphyses (Figure 12). An obvious complication is fracture.

### The Sclerosing Dysplasias

Four skeletal dysplasias are the result of abnormal metaphyseal and periosteal ossification caused by either excessive osteoid production or deficient osteolysis. The primary radiographic feature is increased bone density. They include osteopetrosis, melorheostosis, osteopoikilosis, and osteopathia striata.

Osteopetrosis (Albers-Schönberg disease) is an inherited, autosomal dominant (delayed type) condition presenting radiographically with diffuse bone sclerosis (Figure 13). Its characteristic feature has been described as a “bone within a bone” appearance.<sup>11</sup> These patients are relatively asymptomatic.<sup>1</sup>

Melorheostosis is a rare dysplasia that is not believed to be hereditary.<sup>1</sup> The patient may or may not have associated symptomatology.<sup>9</sup> The radiographic presentation is typically limited to a single limb and consists of hyperos-



Figure 21: Vascular calcification: phleboliths. A, Phleboliths may be solitary or multiple, as demonstrated in this example of the distal leg. B, Numerous phleboliths are seen throughout the calf in this patient with chronic venous insufficiency.



Figure 22: Vascular calcification: Moenckeberg's sclerosis. The classic presentation is shown in this diabetic patient (DP view). Tubular, serpiginous calcifications are easily identified in the first intermetatarsal space following the course of arterial vessels.

tosis along the bone's periphery, extending along its entire length in many cases (Figure 14). This picture simulates wax flowing down the side of a candle and has a wavy, sclerotic bony contour usually along the bone's endosteal surface.



Figure 23: Vascular calcification: atherosclerosis. In contrast to Moenckeberg's sclerosis, the calcifications that follow the course of arterial vessels are patchy and solid.

Osteopoikilosis is an inherited skeletal dysplasia that is not associated with symptomatology. Numerous small, well-defined and homogeneous circular foci of increased density are seen radiographically, mimicking multiple bone islands (Figure 15). The distribution of these lesions is periarticular and symmetric, found at the ends of long bones.<sup>1</sup> They may also be found in tarsal bones.



Figure 19: Generalized calcinosis: tumoral calcinosis. A large periarticular mass demonstrates calcification adjacent to the proximal interphalangeal joint of a toe.



Figure 20: Dystrophic calcification: trauma. This patient related a recent injury to the heel. Multiple areas of ill-defined soft tissue calcification are identified (arrows).

**Osteopoikilosis is characterized by numerous small, well-defined and homogeneous circular foci of increased density.**

Osteopathia striata is most likely inherited and is typically asymptomatic.<sup>9</sup> Radiographically, linear, regular bands of increased density extend from the metaphysis to the diaphysis, running parallel to the shaft (Figure 16).

### Fibrous Dysplasia

Pathologically, normal bone undergoing physiologic resorption is replaced with fibrous tissue.<sup>12</sup> It may occur in one (monostotic) or many (polyostotic) bones. In tubular bones its location is intramedullary and diaphyseal, with only occasional epiphyseal involvement.<sup>1</sup> Radiographically the lesion is somewhat radiolucent with a hazy, ground glass (homogeneous) quality, and may appear expansile with endosteal scalloping (Figure 17).

### Soft Tissue Manifestations of Systemic Disorders

The primary radiographic soft tissue findings are calcification and ossification. Calcification appears as irregular punctate, circular, linear, or plaque-like radiodense areas with no cortical or trabecular structure.<sup>1</sup> Ossification, in contrast, demonstrates a trabecular pattern and a thin, cortex-like periphery.

However, if the lesion is small, the distinction may be impossible to determine visually.

### Calcification

Conditions that lead to soft tissue calcification have been categorized as metastatic calcification, general-

ized calcinosis, and dystrophic calcification.<sup>1</sup> Metastatic calcification (Figure 18) results from a disturbance in calcium or phosphorus metabolism. Examples include hyperparathyroidism, hypoparathyroidism, renal osteodystrophy, hypervitaminosis D, and sarcoidosis. Generalized calcinosis (Figure 19) presents as calcium deposition in the skin and subcutaneous tissue in the presence of normal calcium metabolism. Examples include collagen vascular disorders (scleroderma, dermatomyositis), idiopathic tumoral calcinosis, and idiopathic calcinosis universalis.

Tumoral calcinosis is rare in the foot but has been reported adjacent to a sesamoid<sup>13</sup> and a first-metatarsal-head medial eminence,<sup>14</sup> at the tip of digits,<sup>15</sup> and posterior to the ankle joint. With dystrophic calcification (Figure 20), calcium





Figure 24: Calcific tendonitis. A, Achilles tendon calcification near its enthesis. B, Calcification of extensor tendons along the dorsum of the midfoot.

is deposited in damaged or devitalized tissue in the absence of a generalized metabolic derangement. Examples include neoplasm, inflammation, and trauma. Soft tissue calcification has been associated with local corticosteroid injection of the heel<sup>16</sup> and intra-articular corticosteroid injection of the small joints of the hand.<sup>17</sup> Extensive involvement of the foot suggests sarcoma.<sup>18</sup>

Vascular calcifications are frequently encountered in the foot and leg. Phleboliths, found in veins, are circular or elliptical in shape (Figure 21); they possess a thin calcific outline with a relatively lucent center. Phleboliths may be a solitary finding or multiple. Phleboliths have also been described associated with hemangioma of the foot.<sup>19</sup> Two types of arterial calcification have been described: Moenckeberg's



Figure 25: Soft tissue ossification: Achilles tendon. A, Notice the trabeculations in the lesion, which distinguish it from calcification. B, A much larger ossification; this image also demonstrates Moenckeberg's sclerosis.

sclerosis and atherosclerosis. Moenckeberg's sclerosis (Mönckeberg's arteriosclerosis, medial calcific sclerosis) is calcification of the vessel's tunica media layer. These lesions generally do not obstruct blood flow through the vessel. They are commonly seen in patients with diabetes.

Radiographically, Moenckeberg's sclerosis has a characteristic appearance consisting of dual, curvilinear tubular




Figure 26: Generalized periostitis: venous stasis. A, Irregular, wavy periostitis is identified along the lateral and medial margins of the distal tibia and fibular diaphysis, respectively. B, This patient, with chronic venous insufficiency (AP view), demonstrates solid periostitis along the distal tibial diaphysis with diffuse soft tissue calcification (phleboliths).

calcifications running parallel to one another (Figure 22). They may be serpiginous and follow the anatomic course of the vessel. Not uncommonly the path of calcification is discontinuous. The term atheroscle-




Figure 27: Heterotopic soft tissue ossification. A, Myositis ossificans circumscripta. A lesion identified in the plantar fascia musculature. B, Myositis ossificans associated with neuropathic osteoarthropathy, posterior ankle.

rosis refers to plaque formation along the tunica intima layer of the vessel. Progressive calcification at any particular site can lead to obstruction of the vessel. Radiographically, one notes patchy calcifications in the soft tissues that follow the path of vessels (Figure 23).

The site of soft tissue calcification may provide a clue to the underlying systemic disorder.<sup>1</sup> Periarticular calcification is associated with hyperparathyroidism, hypervitaminosis D, and collagen vascular disease. Calcifications of tendons and bursae are associated with calcium pyrophosphate dihydrate (CPPD) and calci-

um hydroxyapatite crystal deposition disease. Arterial calcification is frequently seen with diabetes mellitus, hypervitaminosis D, and renal osteodystrophy. Calcification of cartilage (chondrocalcinosis) suggests CPPD

deposition and hemorrhage. Sheet-like collections in the lower leg are associated with tissue injury and compartment syndrome.

The term calcific tendinitis (Figure 24) has been used to describe tendon calcification resulting from the deposition of calcium hydroxyapatite

## The site of soft tissue calcification may provide a clue to the underlying systemic disorder.<sup>1</sup>

crystals.<sup>20</sup> Its etiology is closely associated with trauma, but injury is not prerequisite. Pain is the most common associated symptom. Holt and Keats illustrate examples involving the Achilles, peroneus, and forefoot flexor tendons.<sup>20</sup> Similar reports include examples involving the peroneal tendon,<sup>21,22</sup> although they are not identified specifically as calcific tendinitis. Radiographic evidence of

is the most common form (Figure 27A). Several cases in the foot have been described.<sup>29-32</sup>

3) Myositis ossificans associated with neurologic disease (Figure 27B) may be due to a variety of neurologic causes.<sup>7</sup>

4) Pseudomalignant myositis ossificans is similar to the *circumscripta* form but without a history of antecedent trauma.<sup>33</sup>

## The term myositis ossificans is used to describe non-neoplastic heterotopic soft tissue calcification.

adjacent bone erosion, mimicking malignancy, has been reported.<sup>23</sup>

### Ossification

Causes of soft tissue ossification include neoplasm (sarcoma), trauma (myositis ossificans, ossifying hematoma), and venous insufficiency.

Achilles tendon ossification (Figure 25) is frequently associated with pain.<sup>24-26</sup> Interestingly, MRI analysis of several cases failed to demonstrate inflammatory changes, even with acute symptoms.<sup>27</sup> Morris and associates have presented a classification of Achilles tendon lesion based on anatomic location, including calcification and ossification.<sup>28</sup>

Generalized periostitis is associated with venous stasis (Figure 26), hypertrophic osteoarthropathy (primary and secondary), thyroid acropachy, and hypervitaminosis A.<sup>1,11</sup> Radiographically, varying degrees of periosteal new bone formation is possible, but typically it is thick and undulating (irregular).

The term myositis ossificans is used to describe non-neoplastic heterotopic soft tissue calcification. Four categories are described in the literature, relating to underlying etiology (or lack of):

1) Myositis ossificans progressiva is a rare genetic dysplasia with associated congenital osseous abnormalities.<sup>7</sup>

2) Myositis ossificans *circumscripta*, caused by localized trauma,

Yochum and Rowe<sup>34</sup> warn that the early stages of myositis ossificans *circumscripta* may mimic sarcoma. It initially presents as a soft tissue swelling that proceeds to an ill-defined calcific density followed by ossification. A radiolucent zone may be seen between the lesion and adjacent bone, which helps to distinguish it from sarcomatous soft tissue extension. **PM**

### References

<sup>1</sup> Resnick D: Bone and joint imaging, ed 3, Philadelphia, 2004, Saunders.

<sup>2</sup> Bullough PG, Vigoria VJ: Orthopaedic pathology, ed 5, 2009, Mosby.

<sup>3</sup> Christman RA: Misconceptions regarding use of the term "osteoporosis," *J Cur Podiatr Med* 35(10):11, 1986.

<sup>4</sup> Genant HK, Vogler JB, Block JE: Radiology of osteoporosis. In Riggs BL, Melton LJ, editors: *Osteoporosis: etiology, diagnosis, and management*, Philadelphia, 1995, Lippincott Williams & Wilkins.

<sup>5</sup> Jhamaria NL, Lal KB, Udawat M et al: The trabecular pattern of the calcaneum as an index of osteoporosis, *J Bone Joint Surg* 65B(2):195, 1983.

<sup>6</sup> Cockshott WP, Occleshaw CJ, Weber C et al: Can a calcaneal morphologic index determine the degree of osteoporosis? *Skeletal Radiol* 12:119, 1984.

<sup>7</sup> Greenfield GB: Radiology of bone diseases, ed 4, Philadelphia, 1986, Lippincott.

<sup>8</sup> Jaffe HL: Metabolic, degenerative, and inflammatory diseases of bones and joints, Philadelphia, 1972, Lea & Febiger.

<sup>9</sup> Berquist TH: Imaging of the foot and ankle, ed 3, Philadelphia, 2010, Lippincott Williams & Wilkins.

<sup>10</sup> Aegerter E, Kirkpatrick JA: Ortho-

pedic diseases, ed 4, Philadelphia, 1975, Saunders.

<sup>11</sup> Forrester DM, Kricun ME, Kerr R: Imaging of the foot and ankle, Rockville, Md, 1988, Aspen.

<sup>12</sup> Yochum TR, Rowe LJ: Essentials of skeletal radiology, vol 2, ed 3, Baltimore, 2004, Williams & Wilkins.

<sup>13</sup> Slomowitz M, Nixon B, Mott RC: Tumoral calcinosis of the foot: case report and literature review, *JFS* 29(3):278, 1990.

<sup>14</sup> Caspi I, Friedman B, Horoszovski H: Tumoral calcinosis simulating osteomyelitis, *JFS* 28(6):547, 1989.

<sup>15</sup> Black JR, Sladek GD: Tumoral calcinosis in the foot and hand, *J Am Podiatr Assoc* 73(3):153, Mar 1983.

<sup>16</sup> Conti RJ, Shinder M: Soft tissue calcifications induced by local corticosteroid injection, *JFS* 30(1):34, 1991.

<sup>17</sup> Dalinka MK, Stewart V, Bomalaski JS, et al: Periarticular calcifications in association with intra-articular corticosteroid injections, *Radiology* 153:615, 1984.

<sup>18</sup> Wu KK: Tumoral calcinosis with extensive pedal involvement, *JFS* 29(4):388, 1990.

<sup>19</sup> Miller SJ, Patton GW, Xenos D, Wulf MR: Multiple capillary hemangiomas of the foot with associated phleboliths, *J Am Podiatr Assoc* 70(7):364, 1980.

<sup>20</sup> Holt PD, Keats TE: Calcific tendinitis: a review of the usual and unusual, *Skeletal Radiol* 22:1, 1993.

<sup>21</sup> Wirtz PD, Long DH, Vito GR: Calcification within the peroneus brevis tendon, *J Am Podiatr Med Assoc* 77(6):292, 1987.

<sup>22</sup> Lepow GM, Korfin DH: Calcification of an accessory peroneal tendon in an athlete, *J Am Podiatr Med Assoc* 75(6):323, 1985.

<sup>23</sup> Hayes CW, Rosenthal DI, Plata MJ, Hudson TM: Calcific tendinitis in unusual sites associated with cortical bone erosion, *AJR* 149:967, 1987.

<sup>24</sup> Lotke PA: Ossification of the Achilles tendon, *J Bone Joint Surg* 52A(1):157, 1970.

<sup>25</sup> Raynor KJ, McDonald RJ, Edelman RD, Parkinson DE: Ossification of the Achilles tendon, *J Am Podiatr Med Assoc* 76(12):688, 1986.

<sup>26</sup> Hatori M, Kita A, Hashimoto Y et al: Ossification of the Achilles tendon: a case report, *Foot & Ankle* 15(1):44, 1994.

<sup>27</sup> Yu JS, Witte D, Resnick D, Pogue W: Ossification of the Achilles tendon: imaging abnormalities in 12 patients, *Skeletal Radiol* 23:127, 1994.

<sup>28</sup> Morris KL, Giacomelli JA, Granoff D: Classification of radiopaque lesions of the tendo Achillis, *JFS* 29(6):533, 1990.

<sup>29</sup> Kaminsky SL, Corcoran D, Chubb WF, Pulla RJ: Myositis ossificans: pedal manifestations, *JFS* 31(2):173, 1992.

<sup>30</sup> Fuselier CO, Tlapak TA, Sowell RD:

Heterotopic ossification (myositis ossificans) in the foot, *J Am Podiatr Med Assoc* 76(9):524, 1986.

<sup>31</sup> Herring KM, Levine BD: Myositis ossificans of traumatic origin in the foot, *JFS* 31(1):30, 1992.

<sup>32</sup> Mandracchia V, Mahan KT, Pruzansky J, Uricchio JN: Myositis ossificans: a report of a case in the foot, *J Am Podiatr Assoc* 73(1):31, 1983.

<sup>33</sup> Ogilvie-Harris DJ, Hons CB, Fornasier VL: Pseudomalignant myositis ossificans: heterotopic new-bone formation without a history of trauma, *J Bone Joint Surg* 62A(8):1274, 1980.

<sup>34</sup> Yochum TR, Rowe LJ: *Essentials of skeletal radiology*, vol 1, ed 3, Baltimore, 2004, Williams & Wilkins.

**Additional References**

Martinez S: Tumoral calcinosis: 12 years later. *Semin Musculoskelet Radiol* 6: 331, 2002.

Bansal A: The dripping candle wax sign. *Radiology* 246: 638, 2008.

Fotiadou A, Arvaniti M, Kiriakou V, et

al.: Type II autosomal dominant osteopetrosis: radiological features in two families containing five members with asymptomatic and uncomplicated disease. *Skeletal Radiol* 38: 1015, 2009.

Stoker DJ: Osteopetrosis. *Semin Musculoskelet Radiol* 6: 299, 2002.

Williams HJ, Davies AM and Chapman S: Bone within a bone. *Clinical Radiology* 59: 132, 2004.

Berry JL, Davies M and Mee AP: Vitamin D metabolism, rickets, and osteomalacia. *Semin Musculoskelet Radiol* 6: 173, 2002.

Morselli LL, Manetti L, Cosci C, et al.: Bone and joint alterations in acromegaly. *Journal of Orthopaedics and Traumatology* 7: 169, 2006.

Mughal Z: Rickets in childhood. *Semin Musculoskelet Radiol* 6: 183, 2002.

Pitt MJ, Morgan SL, Lopez-Ben R, et al.: Association of the presence of bone bars on radiographs and low bone mineral density. *Skeletal Radiol* 40: 905, 2011.

Quek ST and Peh WC: Radiology of

osteoporosis. *Semin Musculoskelet Radiol* 6: 197, 2002.

Whitehouse RW: Paget's disease of bone. *Semin Musculoskelet Radiol* 6: 313, 2002.

Wilson LC and Hall CM: Albright's hereditary osteodystrophy and pseudohypoparathyroidism. *Semin Musculoskelet Radiol* 6: 273, 2002.



**Dr. Christman** recently retired. He was previously a tenured professor at Western University of Health Sciences' College of Podiatric Medicine in Pomona, CA and, prior to that, Director of Radiology at Temple University's School of Podiatric Medicine in Philadelphia, PA. He is the editor and author of the textbook *Foot and Ankle Radiology*; its 2nd edition was published in 2015.

**CME EXAMINATION**

1) Osseous resorption in cortical bone may appear as:

- A) band-like/transverse radiolucent areas in the metaphysis.
- B) subchondral spotty radiolucent areas.
- C) intracortical tunneling.
- D) diffuse, homogeneous "coarsening" of trabeculae.

2) A major cause of diffuse osteopenia is:

- A) osteoporosis.
- B) osteoarthritis.
- C) osteosarcoma.
- D) osteopoikilosis.

3) Generalized subperiosteal bone resorption is characteristic of:

- A) pseudohypoparathyroidism.
- B) hyperparathyroidism.
- C) fluorosis.
- D) hypervitaminosis A.

4) Generalized soft tissue thickening, increased girth of metatarsals, decreased girth of proximal phalanges, and joint space widening are best associated with:

- A) hyperparathyroidism.
- B) pseudohypoparathyroidism.
- C) acromegaly.
- D) hypothyroidism.

5) Short metatarsals or phalanges are best associated with:

- A) hypothyroidism.
- B) pseudohypoparathyroidism.
- C) hyperparathyroidism.
- D) hypervitaminosis A.

6) The following is NOT one of the three classes of osteoporosis:

- A) generalized.
- B) regional.
- C) localized.
- D) field.

- 7) A radiographic finding seen with reflex sympathetic dystrophy is:
- A) spotty loss of bone density.
  - B) intracortical tunneling.
  - C) coarse trabeculations.
  - D) cortical thinning.
- 8) A major cause of regional osteoporosis in the lower extremity is:
- A) swimming.
  - B) running.
  - C) walking.
  - D) immobilization.
- 9) A radiographic feature of rickets is:
- A) straight, narrow tubular bone.
  - B) sclerotic epiphysis.
  - C) narrowing of the metaphysis.
  - D) irregular outline or zone of provisional calcification.
- 10) Hypovitaminosis C is also known as:
- A) rickets.
  - B) scurvy.
  - C) lead poisoning.
  - D) hypervitaminosis D.

The author(s) certify that they have NO affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest), or non-financial interest (such as personal or professional relationships, affiliations, knowledge, or beliefs) in the subject matter or materials discussed in this manuscript.

## PM's CME Program

Welcome to the innovative Continuing Education Program brought to you by *Podiatry Management Magazine*. Our journal has been approved as a sponsor of Continuing Medical Education by the Council on Podiatric Medical Education.

### Now it's even easier and more convenient to enroll in PM's CE program!

You can now enroll at any time during the year and submit eligible exams at any time during your enrollment period.

### CME articles and examination questions from past issues of *Podiatry Management* can be found on the Internet at

**podiatry.com/cme**. Each lesson is approved for 1.5 hours continuing education contact hours. Please read the testing, grading and payment instructions to decide which method of participation is best for you.

Please call 516-521-4474 if you have any questions. A personal operator will be happy to assist you.

Each of the 10 lessons will count as 1.5 credits; thus a maximum of 15 CME credits may be earned during any 12-month period. You may select any 10 in a 24-month period.

***The Podiatry Management Magazine CME Program is approved by the Council on Podiatric Medical Education as a provider of continuing education in podiatric medicine. Podiatry Management Magazine CME has approved this activity for a maximum of 1.5 Continuing Education Contact Hours for each exam successfully completed.***

PM's privacy policy can be found at [podiatry.com/privacy.cfm](http://podiatry.com/privacy.cfm).

This CME is valid for CPME-approved credits for three (3) years from the date of publication.

# Enrollment/Testing Information and Answer Sheet

**Note:** If you are mailing your answer sheet, you must complete all info. on the front and back of this page and mail with your credit card information to: **Program Management Services, 12 Bayberry Street, Hopewell Junction, NY 12533.**

## TESTING, GRADING AND PAYMENT INSTRUCTIONS

(1) Each participant achieving a passing grade of 70% or higher on any examination will receive an official computer form stating the number of CE credits earned. This form should be safeguarded and may be used as documentation of credits earned.

(2) Participants receiving a failing grade on any exam will be notified and permitted to take one re-examination at no extra cost.

(3) All answers should be recorded on the answer form below. For each question, decide which choice is the best answer, and circle the letter representing your choice.

(4) Complete all other information on the front and back of this page.

(5) Choose one out of the 3 options for testgrading: mail-in, fax, or phone. To select the type of service that best suits your needs, please read the following section, "Test Grading Options".

## TEST GRADING OPTIONS

### Mail-In Grading

To receive your CME certificate, complete all information and mail with your credit card information to: **Program Management Services, 12 Bayberry Street, Hopewell Junction, NY 12533.**

**PLEASE DO NOT SEND WITH SIGNATURE REQUIRED, AS THESE WILL NOT BE ACCEPTED.**

There is **no charge** for the mail-in service if you have already enrolled in the annual exam CME program, and we receive this exam during your current enrollment period. If you are not enrolled, please send \$35.00 per exam, or \$299 to cover all 10 exams (thus saving \$51 over the cost of 10 individual exam fees).

### Facsimile Grading

To receive your CME certificate, complete all information and fax 24 hours a day to 1631-532-1964. Your test will be dated upon receipt and a PDF of your certificate of completion will be sent to the Email address on file with us. Please allow 5 business days for the return of your certificate. This service is available for \$2.95 per exam if you are currently enrolled in the 10-exam CME program, and can be charged to your Visa, MasterCard, or American Express.

If you are *not* enrolled in the 10-exam CME program, the fee is \$35 per exam.

### Phone-In Grading

You may also complete your exam by using the toll-free service. Call 516-521-4474 from 10 a.m. to 5 p.m. EST, Monday through Friday. Your CME certificate will be dated the same day you call and mailed within 48 hours. There is a \$2.95 charge for this service if you are currently enrolled in the 10-exam CME program, and this fee can be charged to your Visa, Mastercard, American Express, or Discover. If you are not currently enrolled, the fee is \$35 per exam. When you call, please have ready:

1. Program number (Month and Year)
2. The answers to the test
3. Credit card information

In the event you require additional CME information, please contact PMS, Inc., at **516-521-4474**.

## ENROLLMENT FORM & ANSWER SHEET

*Please print clearly...Certificate will be issued from information below.*

Name \_\_\_\_\_ Email Address \_\_\_\_\_

Please Print:      FIRST                      MI                      LAST

Address \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Charge to:  Visa     MasterCard     American Express

Card # \_\_\_\_\_ Exp. Date \_\_\_\_\_ Zip for credit card \_\_\_\_\_

**Note: Credit card is the only method of payment. Checks are no longer accepted.**

Signature \_\_\_\_\_ Email Address \_\_\_\_\_ Daytime Phone \_\_\_\_\_

State License(s) \_\_\_\_\_ Is this a new address? Yes \_\_\_\_\_ No \_\_\_\_\_

**Check one:**  I am currently enrolled. (If faxing or phoning in your answer form please note that \$2.95 will be charged to your credit card.)

I am not enrolled. Enclosed is my credit card information. Please charge my credit card \$35.00 for each exam submitted. (plus \$2.95 for each exam if submitting by fax or phone).

I am not enrolled and I wish to enroll for 10 courses at \$299.00 (thus saving me \$51 over the cost of 10 individual exam fees). I understand there will be an additional fee of \$2.95 for any exam I wish to submit via fax or phone.

*Over, please*

**EXAM #5/24**  
**Radiology of Systemic Disorders**  
**Affecting the Foot and Ankle**  
**(Christman)**

Circle:

- |            |             |
|------------|-------------|
| 1. A B C D | 6. A B C D  |
| 2. A B C D | 7. A B C D  |
| 3. A B C D | 8. A B C D  |
| 4. A B C D | 9. A B C D  |
| 5. A B C D | 10. A B C D |

**Medical Education Lesson Evaluation**

Strongly agree [5]	Agree [4]	Neutral [3]	Disagree [2]	Strongly disagree [1]
--------------------------	--------------	----------------	-----------------	-----------------------------

- 1) This CME lesson was helpful to my practice \_\_\_\_
- 2) The educational objectives were accomplished \_\_\_\_
- 3) I will apply the knowledge I learned from this lesson \_\_\_\_
- 4) I will make changes in my practice behavior based on this lesson \_\_\_\_
- 5) This lesson presented quality information with adequate current references \_\_\_\_
- 6) What overall grade would you assign this lesson?  
A B C D
- 7) This activity was balanced and free of commercial bias.  
Yes \_\_\_\_ No \_\_\_\_
- 8) What overall grade would you assign to the overall management of this activity?  
A B C D

How long did it take you to complete this lesson?  
\_\_\_\_hour \_\_\_\_minutes

What topics would you like to see in future CME lessons?  
Please list :

---

---

---

---

---